

# Thompson Sampling for Infinite-Horizon Discounted Decision Processes

Daniel Adelman<sup>1</sup>, Cagla Keceli<sup>1</sup> and Alba V. Olivares Nadal<sup>2</sup>

<sup>1</sup>Booth School of Business, The University of Chicago, Chicago, IL 60637

<sup>2</sup>UNSW Business School, University of New South Wales, Sydney, Australia

## Abstract

We model a Markov decision process, parametrized by an unknown parameter, and study the asymptotic behavior of a sampling-based algorithm, called Thompson sampling. The standard definition of regret is not always suitable to evaluate a policy, especially when the underlying chain structure is general. We show that the standard (expected) regret can grow (super-)linearly and fails to capture the notion of learning in realistic settings with non-trivial state evolution. By decomposing the standard (expected) regret, we develop a new metric, called the expected residual regret, which forgets the immutable consequences of past actions. Instead, it measures regret against the optimal reward moving forward from the current period. We show that the expected residual regret of the Thompson sampling algorithm is upper bounded by a term which converges exponentially fast to 0. We present conditions under which the posterior sampling error of Thompson sampling converges to 0 almost surely. We then introduce the probabilistic version of the expected residual regret and present conditions under which it converges to 0 almost surely. Thus, we provide a viable concept of learning for sampling algorithms which will serve useful in broader settings than had been considered previously.

---

# 1 Introduction

We consider a control problem in which a decision-maker (DM) interacts with an environment without knowing the value of a parameter encoded into this environment. This environment is modeled by a discrete-time Markov decision process (MDP). The DM applies a control and, in return, the system moves to the next state and generates a reward. Neither of the outputs are deterministic; they are contingent on the current state, control and value of the unknown parameter. Using the state transition and reward information, the DM picks a control which yields the subsequent transition and reward.

Our model is mainly shaped by the stochastic adaptive control literature and, in particular, complements [Kim \(2017\)](#). Numerous prominent papers parametrize MDP's by an unknown parameter and estimate this parameter using different methods. We investigate the asymptotic performance of a popular parameter estimation algorithm, Thompson sampling (TS), first described by [Thompson \(1933\)](#). Aiming to maximize the total reward, yet not knowing the underlying parameter that determines the rewards and state transitions, the DM faces a trade-off between choosing controls tied to unexplored parameter candidates and choosing controls which are likely to yield high rewards. This exploration vs. exploitation trade-off is often addressed in online decision-making problems by TS. After estimating the unknown parameter by TS, the DM selects a control while assuming the estimate is the true parameter.

Despite the logarithmic expected regret guarantees in bandit settings ([Agrawal and Goyal 2012](#)), we document through examples that the expected regret no longer has logarithmic guarantees when the chain structure underneath the MDP is generic. In particular, we document how the expected regret can grow (super-)linearly. This means that the standard notion of expected regret used in the literature, i.e., the gap between the cumulative reward of the DM and the cumulative reward of an omniscient agent, is not fully informative when the underlying chain is general. The expected regret is unable to discern that learning is taking place, even though it is. These kinds of examples may be implicitly known to researchers who work in this area, yet they have not been documented in the literature to the best of our knowledge.

We provide the first theoretical results that capture a different notion of (expected) regret of the TS algorithm. In particular, we extend a result of [Kim \(2017\)](#) (on the convergence of the expected posterior error) to discounted-reward MDP's and to general state and control spaces. We assume Borel state-control spaces, allowing the spaces to be infinite. Thus, our work allows questions of

---

regret and sampling algorithms to be addressed in broader settings in which they have not been addressed before. Extending the performance guarantees of the new notion of regret to MDP’s with a general chain structure is an important potential avenue of study.

The literature on parametrized MDP’s almost always considers the long-run average regret per period. In contrast, we study guarantees when the performance criterion is the expected infinite-horizon discounted reward. Since problems of economic significance are often most properly formulated with discounting, having the objective function formulated as an infinite-horizon discounted reward problem would allow the ideas of learning through sampling to be applied in these settings.

We develop a general canonical probability space for adaptive learning algorithms based on sampling, of which TS is a special case. [Kim \(2017\)](#) and [Banjević and Kim \(2019\)](#) construct a similar probability space for TS, but do not include the sampled parameters in the sample space. We are not aware of other work which incorporates this canonical formulation. Having the sampling algorithms posed in a coherent framework allows for it to be understood and studied rigorously. For example, the formulation makes evident that the underlying process is history-dependent, i.e., not Markovian. To make this clear, we will refer to the adaptive version of the process with an unknown parameter as  $\theta$ -MDP instead of MDP.

We offer a novel decomposition of the standard notion of expected regret. Our notion of expected regret can be computed at any finite period  $n$ , from the vantage point of a DM in period 0. We identify three distinct components of the expected regret, only one of which is always suitable for learning problems. The first component is the expected regret associated with the past, i.e., from period 0 up to period  $n$ . This quantifies the difference between the expected reward received by the DM and the expected reward of the policy which knows the true parameter. The second component is the expected regret that emanates from the policy’s state in period  $n$ , which is inextricably tied to the past decisions. It captures the expected infinite-horizon (future) consequences of being in a suboptimal state in period  $n$ . The third and final component captures the DM’s ability to implement what is optimal moving forward, conditional on the period- $n$  state. This component is the only component that is “controllable” by the DM, once they have arrived at the period- $n$  state. It reflects how the best opportunities available to the DM from period  $n$  onward compare to how the system will evolve under the sampling policy. Our analysis focuses on this concept which we call *expected residual regret*. The expected residual regret forgets both the past and the future “sunk” portions of the total regret.

Decomposing the standard expected regret is a novel approach to view and quantify the per-

---

formance of a policy. Through this decomposition, we connect the notion of expected regret to a concept called *asymptotic discount optimality* (ADO) (Schäl 1987, Hernández-Lerma 2012). We demonstrate how the ADO concept relates to expected residual regret. To the best of our knowledge, this concept has not been used within the sampling context before and has not been connected to the notion of (expected) regret.

We show that the posterior belief on the true parameter converges to 1 when TS is implemented; this behavior is called *complete learning*. Complete learning is a stronger result than learning in expectation, i.e., expected posterior belief, because it gives assurance about the individual sample paths. To the best of our knowledge, most results are on the expected-valued version of posterior belief. The notion of complete learning has been studied frequently in bandit contexts, including the prominent work of Freedman (1963), yet to our knowledge it has not been applied to Borel state-control spaces and has not been analyzed under the discounted reward criterion. In the adaptive learning literature, there is an alternative to TS, known as the minimum contrast estimator, which is shown to achieve complete learning (Hernández-Lerma 2012). However, complete learning has not been connected to the concept of ADO, and there is no notion of probabilistic ADO.

Building on expected residual regret, we define its sample-path version, called *probabilistic residual regret* or *residual regret*. Similar to expected residual regret, it captures the DM’s ability to implement what is optimal moving forward, but starting from the random state of the MC in period  $n$ . It quantifies the difference in optimal rewards and TS-driven rewards. Therefore, the probabilistic residual regret is a random quantity whose expected value is the expected residual regret. Conceptually, probabilistic residual regret is similar to complete learning as they are both concerned with individual sample paths.

The paper is organized as follows. In Section 2, we provide additional literature review. Then we motivate the need to modify the definition of expected regret since the standard notion of expected regret can grow linearly in non-trivial settings. In Section 3, we model the sampling algorithm and formulate the canonical probability space which involves the sampled parameters. In Section 4, we characterize TS by defining the posterior update and control selection mechanisms. In Section 5, we decompose the standard notion of expected regret into components and interpret each component, highlighting expected residual regret as the only “actionable” component. In Section 6, we provide an asymptotic analysis of the expected residual regret. In Section 7, we show that TS exhibits complete learning, i.e., the posterior sampling error converges to 0 almost surely. We also show that probabilistic residual regret converges to 0 almost surely.

---

## 2 Literature Review and Motivation

Here, we highlight papers of similar setup and provide an example where the standard expected regret fails to discern learning.

### 2.1 Literature Review

TS has proved not only successful for multi-armed bandit problems (MAB) (a degenerate case of MDP); but also for parametrized MDP's, i.e.,  $\theta$ -MDP's, which need not be Markovian. The work on  $\theta$ -MDP's typically requires assumptions on the underlying chain to give performance guarantees on the expected regret. For example, [Kim \(2017\)](#) shows that TS achieves asymptotically optimal expected regret when the Markov chain under the optimal policy, which knows the true parameter, is ergodic.<sup>1</sup> Another example is [Gopalan and Mannor \(2015\)](#), which provides a probabilistic logarithmic upper bound on the expected regret of TS, assuming that the starting state is recurrent under the optimal policy generated by any of the possible parameters. Although [Kearns and Singh \(2002\)](#) and [Leike et al. \(2016\)](#) assume a more general chain setting, they study decision processes over finite state-control spaces. In contrast to the finite state-control spaces or the expected average-reward criterion of these works, we assume a general state-control space, under the discounted-reward criterion, similar to [Schäl \(1987\)](#). We impose additional assumptions on the underlying chain to be able to extend the work of [Kim \(2017\)](#) into our more general setting.

[Kim \(2017\)](#) shows that the expected posterior sampling error decays exponentially. We extend this result to a broader framework. We also show that the probabilistic version of the posterior sampling error decays exponentially, which we believe has not been explored yet.

Our work mostly complements [Kim \(2017\)](#) and [Banjević and Kim \(2019\)](#). Both works show good performance guarantees for Thompson sampling under parameter uncertainty, yet our work has a stronger connection with the former since we also assume a finite parameter space. The latter adopts a continuous parameter space, similar to [Hernández-Lerma \(2012\)](#), which results in a setting that is harder to analyze.

The stream of work that analyzes the behavior of TS do not usually consider infinite-horizon, and are most often in the MAB setting, with history-independent samples to the best of our knowledge. [Kalkanli and Ozgur \(2020\)](#) analyze the asymptotic behavior of TS with history-dependence, but in the context of the MAB problem. Although we utilize different methodologies than theirs, our work

---

<sup>1</sup> A Markov chain is *ergodic* if the transition matrix corresponding to every deterministic stationary policy consists of a single recurrent class ([Puterman 2014](#)).

---

carries this analysis into the under-explored context of the  $\theta$ -MDP.

## 2.2 Motivation

TS has been shown to have good performance in the MAB setting. The MAB problem is equivalent to a  $\theta$ -MDP with no state, or alternatively, a one-step  $\theta$ -MDP. In each decision period  $t$ , the DM samples a parameter from the posterior distribution. By treating the sample as the true parameter, the DM chooses a control, i.e., plays one of the constantly-many arms, and immediately observes a reward. The reward of each arm is generated according to some fixed (unknown) distribution and the objective is to maximize the total expected reward. Arms' rewards are generated independently of each other. Let  $\mu_i$  denote the (unknown) expected reward of arm  $i$  and  $i(t)$  be the arm played in period  $t$ . The expected finite-time regret is the expected total difference between the optimal strategy of pulling the arm with the highest mean and the strategy followed by the DM, i.e.,

$$E[\text{Regret}(n)] := E \left[ \sum_{t=1}^n (\mu^* - \mu_{i(t)}) \right].$$

where  $\mu^* := \max_i \mu_i$ . In the pioneering [Lai and Robbins \(1985\)](#), the (expected) regret of any bandit algorithm is lower bounded, in the limit, by

$$E[\text{Regret}(n)] \geq \left[ \sum_{i=2}^K \frac{\Delta_i}{D(\mu_i || \mu^*)} + o(1) \right] \ln(n), \tag{1}$$

where  $D$  denotes the Kullback–Leibler divergence and  $\Delta_i := \mu^* - \mu_i$ . The bound in (1) shows that the best achievable expected regret is of order  $\ln(n)$ . Complementing the logarithmic lower bound for any bandit algorithm, [Agrawal and Goyal \(2012\)](#) upper bounds the expected regret of the TS algorithm by

$$E[\text{Regret}(n)] \leq O \left( \left( \sum_{i=2}^K \frac{1}{\Delta_i^2} \right)^2 \ln(n) \right).$$

Since the order of the upper bound on the expected regret of TS matches the logarithmic lower bound for any algorithm in (1), the expected regret of TS grows logarithmically. We underline that these results are valid for the MAB setting. These results can also apply to the  $\theta$ -MDP setting with a trivial state process, such that the stochastic process is driven by an iid state process<sup>2</sup>. In addition, they assume no discounting of the rewards. Our setting is fundamentally different, i.e., the states evolve based on the controls and are not identically distributed, hence these results do not apply in general. [Example 2.1](#) illustrates a simple yet non-trivial state process.

---

<sup>2</sup> The states evolve independently of the controls and have the same probability distribution.

---

**Example 2.1** (Expected regret grows linearly). Consider the three-state process shown in Figure 1. In period  $t = 0$ , the DM lies in  $x_0$  and can choose either control  $A$  or  $B$ , with an immediate reward of 0. Control  $A$  leads to the state  $x_A$ , and from then onward only control  $A$  can be chosen; the DM is “stuck”. Control  $B$  leads to the state  $x_B$ , and similarly, only control  $B$  can be chosen from then onward. The true parameter can either be  $A$  or  $B$ . We represent the one-step reward generating function by  $R^A(\cdot)$  when  $A$  is the true parameter, and  $R^B(\cdot)$  otherwise. We assume that the prior belief on the true parameter is not degenerate, i.e., not equal to 1.




Figure 1: Constant reward depending only on the first control, picked at  $t=0$ .

In Example 2.1 if the first guess is right, the DM receives a reward of 1 forever and otherwise receives no reward at all. Thus, unlike in the setting of Agrawal and Goyal (2012), in broader settings the expected finite-horizon regret does not necessarily grow logarithmically.

Consider Example 2.1 with an alternative setup such that when the guess at  $t = 0$  is correct, the reward generated in the corresponding state is equal to the number of periods the policy has spent in that state. If the DM makes a wrong guess they are stuck with 0 reward forever, while the oracle earns a sequence of increasing rewards. Here, the expected (undiscounted) regret grows super-linearly. While the standard expected regret grows linearly or super-linearly, nonetheless learning still happens. If the DM receives a reward of 1 in the next period, then they immediately learn whether  $A$  or  $B$  drives the reward process. This example shows the motivation to construct an alternative, more “lenient”, notion of expected regret that forgets the immutable consequences of past actions.

---

### 3 Model Setup

We study a discounted-reward stochastic control problem over an infinite horizon. The underlying MDP is indexed by some parameter  $\theta$ . In Section 3.1, we formulate the MDP when the DM knows  $\theta$ , building on the mathematical framework of [Hernández-Lerma \(2012\)](#). In the subsequent sections, we assume the DM solves the  $\theta$ -MDP using estimates of  $\theta$ , i.e., without knowing the value of  $\theta$ . We construct the probability space in Section 3.2. This lays the groundwork for TS.

#### 3.1 Markov Decision Process for Known $\theta$

Consider a discrete-time MDP,  $(\mathcal{X}, \mathcal{U}, \{\mathcal{U}(x), x \in \mathcal{X}\}, f^\theta, q^\theta)$ . In our setting, the reward and state transition densities depend on the parameter  $\theta \in \mathcal{P}$ , which is a single point in the finite parameter space  $\mathcal{P}$ . Later, when  $\theta$  is assumed to be unknown, we will represent the random sample drawn in period  $t$  by  $\Theta_t$  and its realization by  $\theta_t$ . We reserve upper-case letters to denote random variables, lower-case letters to realizations of random variables, script letters to spaces and sometimes sets, and bold upper-case letters to elements of  $\sigma$ -algebras to be defined. Let  $\mathbb{R}$  denote the set of real numbers, and  $\mathcal{B}(\cdot)$  the Borel  $\sigma$ -algebra of a topological space.

The tuple  $(\mathcal{X}, \mathcal{U}, \{\mathcal{U}(x), x \in \mathcal{X}\}, f^\theta, q^\theta)$  consists of:

1. State space  $\mathcal{X}$ , a Borel space. We denote the system state in period  $t \in \{0, 1, 2, \dots\}$  by  $X_t \in \mathcal{X}$ .
2. Control space  $\mathcal{U}$ , a Borel space. The control applied in period  $t$  is  $U_t \in \mathcal{U}$ .
3. Set of admissible controls  $\mathcal{U}(x)$ , which is a compact subset of  $\mathcal{U}$  for every  $x \in \mathcal{X}$ . Let  $\mathcal{U} = \bigcup_{x \in \mathcal{X}} \mathcal{U}(x)$ . The set of admissible state-control pairs,

$$\mathbb{K} := \{(x, u) \mid x \in \mathcal{X}, u \in \mathcal{U}(x)\},$$

is assumed to be a measurable subset of the product space  $\mathcal{X} \times \mathcal{U}$ .

4. Given  $x_t$  and  $u_t$  in period  $t$ , the system generates random reward  $R_t \in \mathcal{R}_c \subset \mathbb{R}_+$ , where  $\mathcal{R}_c$  is compact and measurable according to a conditional distribution  $F^\theta(\cdot \mid x_t, u_t)$ . The conditional distribution  $F^\theta(\cdot \mid x, u)$  admits a measurable, continuous, one-step reward density<sup>3</sup>  $f^\theta : \mathcal{R}_c \rightarrow \mathbb{R}_+$  with respect to a  $\sigma$ -finite measure  $\lambda$  on  $(\mathcal{R}_c, \mathcal{B}(\mathcal{R}_c))$ , such that

$$F^\theta(\mathbf{R} \mid x, u) := \int_{\mathbf{R}} f^\theta(r \mid x_t = x, u_t = u) d\lambda(r), \quad \forall \mathbf{R} \in \mathcal{B}(\mathcal{R}_c), (x, u) \in \mathbb{K}.$$

---

<sup>3</sup> Radon-Nikodym derivative of  $F^\theta$  with respect to  $\lambda$ .

The expected reward in period  $t$  is

$$r^\theta(x_t, u_t) := \int_{\mathcal{R}_c} r f^\theta(r | x_t, u_t) d\lambda(r), \quad \forall (x_t, u_t) \in \mathbb{K}. \quad (2)$$

5. Given  $x_t$  and  $u_t$  in period  $t$ , the system transitions into random state  $X_{t+1}$  according to a conditional distribution  $Q^\theta(\cdot | x_t, u_t)$ . The conditional distribution  $Q^\theta(\cdot | x, u)$  admits a one-step transition density<sup>4</sup>  $q^\theta : \mathbb{K} \rightarrow \mathcal{X}$  with respect to a  $\sigma$ -finite measure  $\eta$  on  $(\mathcal{X}, \mathcal{B}(\mathcal{X}))$  such that

$$Q^\theta(\mathbf{X} | x, u) := \int_{\mathbf{X}} q^\theta(y | x_t = x, u_t = u) d\eta(y), \quad \forall \mathbf{X} \in \mathcal{B}(\mathcal{X}).$$

Similarly, given  $x_t$  and  $u_t$  in period  $t$ ,

$$Q^\theta(\mathbf{X} | x, u) := \text{Prob}(X_{t+1} \in \mathbf{X} | x_t = x, u_t = u), \quad \forall \mathbf{X} \in \mathcal{B}(\mathcal{X}).$$

**Remark.** Since  $f^\theta(\cdot | x_t, u_t)$  is continuous on compact set  $\mathcal{R}_c$ , it attains its minimum and maximum in  $\mathcal{R}_c$ . Thus, there exists a constant  $M \geq 0$  such that  $|r^\theta(x_t, u_t)| \leq M \forall (x_t, u_t) \in \mathbb{K}$ . To see why this holds, note that

$$\begin{aligned} |r^\theta(x_t, u_t)| &\leq \int_{\mathcal{R}_c} |r f^\theta(r | x_t, u_t)| d\lambda(r) \leq \int_{\mathcal{R}_c} \max(|r f^\theta(r | x_t, u_t)|) d\lambda(r) \\ &= \max_{r \in \mathcal{R}_c} |r f^\theta(r | x_t, u_t)| \lambda(\mathcal{R}_c) < \infty, \end{aligned}$$

where the first inequality follows by (2) and Jensen's inequality, and the equality holds since  $\max(|r f^\theta(r | x_t, u_t)|)$  is constant in  $r$ . Then,  $\max_{r \in \mathcal{R}_c} |r f^\theta(r | x_t, u_t)|$  is finite because  $f^\theta(\cdot | x_t, u_t)$  is continuous on a compact set, and  $\lambda(\mathcal{R}_c)$  is finite since  $\mathcal{R}_c \subset \mathbb{R}_+$  is compact and  $\lambda$  is a  $\sigma$ -finite measure.

The DM aims to maximize the infinite-horizon expected total discounted reward. The *optimal value function*  $\nu^\theta(x)$  represents the maximum such reward starting from state  $x$ , and it depends on the true parameter  $\theta$ . It is known from [Hernández-Lerma and Lasserre \(2012\)](#) that this problem is solved by the Bellman equation in (3).

$$\nu^\theta(x) := \sup_{u \in \mathcal{U}(x)} \left\{ r^\theta(x, u) + \beta \int_{y \in \mathcal{X}} \nu^\theta(y) dQ^\theta(dy | x, u) \right\}, \quad \forall x \in \mathcal{X}, \theta \in \mathcal{P}, \quad (3)$$

where  $\beta \in [0, 1)$  is the discounting factor. If  $\nu^\theta(x)$  is a solution to (3), let  $\mu^\theta$  denote the corresponding optimal policy for parameter  $\theta$ . Under sufficient technical conditions,  $\mu^\theta$  is a stationary, deterministic, and Markovian policy. As showing these conditions holds in this setting are outside of our scope, we make the following assumption:

---

<sup>4</sup> Radon-Nikodym derivative of  $Q^\theta$  with respect to  $\eta$

---

*Assumption 0.* For all  $\theta \in \mathcal{P}$ , there exists a unique solution to (3) such that the supremum is attained, and there exists an optimal policy  $\mu^\theta$  that is stationary, deterministic, and Markovian.

**Remark.** The condition in Assumption 0 is necessary for all of the main results of the paper. Therefore, we will not specifically refer to it in the Lemma, Proposition and Theorem statements.

In this section, we formulated an MDP parametrized by a known parameter  $\theta$ . In Section 3.2, we define the optimal policy in the adaptive setting, i.e., where  $\theta$  is an unknown (fixed) parameter.

### 3.2 Adaptive Learning with Sampling

From this point forward we take  $\theta$  to be an unknown and fixed parameter. A control problem parametrized by an unknown parameter is called an *adaptive* control problem, we refer the reader to [Hernández-Lerma \(2012\)](#) for a comprehensive definition. As defined in Section 3.1, the reward density and transition density are parametrized by the (finitely-many) parameter values. Since we reserve  $\theta$  for the true parameter,  $f^\gamma(\cdot | x, u)$  and  $q^\gamma(\cdot | x, u)$  represent the reward and transition density of an arbitrary parameter  $\gamma \in \mathcal{P}$ . Since  $\theta$  is the underlying true parameter,  $f^\theta(\cdot | x, u)$  and  $q^\theta(\cdot | x, u)$  represent the reward and transition density which drive the actual (observed) process, i.e., the observed reward and the observed next state.

Next, we estimate the unknown parameter  $\theta$  using a generic sampling algorithm.

#### 3.2.1 Overview of Sampling Algorithm.

Had  $\theta$  been known to the DM, the DM would have maximized the objective function by solving the optimization problem in (3) for known  $\theta$ , under Assumption 0. Given the parameter uncertainty, the problem evolves into a learning problem, where the DM needs to choose suitable controls while gathering information on  $\theta$ . To specify how the DM chooses controls in each period  $t$ , we first define the data available at the beginning of each period  $t$ , i.e., the history. We will use history and admissible history interchangeably. In contrast to the setting with known parameter  $\theta$ , when  $\theta$  is unknown the learning procedure makes use of the entire history, i.e., is not Markovian.

Figure 2 illustrates the random variables which drive the learning process, ordered in the sequence of occurrence. The DM observes the history information from period 0 up to period  $t$ , denoted by  $H_t$ . The history vector contains all information up to and including  $X_t$ , i.e., the state in period  $t$ . Provided by the system designer, the function  $\pi_t(\cdot | H_t)$  takes  $H_t$  as input and returns the distribution on the period- $t$  sample  $\Theta_t$ . Although  $\pi_t(\cdot | H_t)$  is a deterministic function of  $H_t$ ,




Figure 2: Evolution of the stochastic process, in the case when  $\theta$  is not known.

it is random due to random  $H_t$ . Next, the DM draws a sample  $\Theta_t$  from the parameter pool, with probability equal to  $\pi_t(\Theta_t | H_t)$ . The randomness of  $\Theta_t$  arises from two sources: due to its dependence on random  $H_t$  through  $\pi_t(\cdot | H_t)$ , and the random nature of sampling. Given the history, the DM solves an optimization problem yielding the “optimal” control  $U_t \in \mathcal{U}(X_t)$ , assuming the true  $\theta$  equals  $\theta_t$ , the realized sample. The history includes only admissible controls, defined in Section 3.1. The state-control pair  $(X_t, U_t)$  gives rise to a random reward  $R_t$ , which depends on  $\theta$ .  $R_t$  is drawn from the reward density  $f^\theta(\cdot | X_t, U_t)$ . Finally, given the transition density  $q^\theta(\cdot | X_t, U_t)$  and the current state-control pair, the system transitions into a random next state  $X_{t+1}$ .

The collection of  $H_t$ ,  $\Theta_t$ ,  $U_t$ ,  $R_t$  and  $X_{t+1}$  constitutes  $H_{t+1}$ . The system designer provides  $f^\gamma(\cdot | x, u)$  and  $q^\gamma(\cdot | x, u)$  for all candidate values  $\gamma \in \mathcal{P}$ , which are utilized to update the belief vector, to be defined in Section 4. Nevertheless, the evolution of the process depends only on  $\theta$ , i.e., nature acts according to  $f^\theta(\cdot | x, u)$  and  $q^\theta(\cdot | x, u)$ . After observing the rewards and transitions in period  $t$ , the DM updates their belief vector  $\pi_{t+1}(\cdot | H_{t+1})$  in period  $t + 1$ . Conditional on the state-control pair, the reward and transitions are mutually independent.<sup>5</sup>

### 3.2.2 Canonical Formulation.

The space of admissible histories up to period  $t$  is denoted by  $\mathcal{H}_t$ . For  $t = 0$ , we have  $\mathcal{H}_0 := \mathcal{X}$ , i.e.,  $H_0 := X_0$ . For  $t \geq 1$ , the space of admissible histories has product form,

$$\mathcal{H}_t := (\mathcal{X} \times \mathcal{P} \times \mathcal{U} \times \mathcal{R}_c)^t \times \mathcal{X} = \mathcal{H}_{t-1} \times \mathcal{P} \times \mathcal{U} \times \mathcal{R}_c \times \mathcal{X}.$$

The history spaces  $\mathcal{H}_0$  and  $\mathcal{H}_t$  ( $t = 1, 2, \dots$ ) are endowed with their product Borel  $\sigma$ -algebras  $\mathcal{B}(\mathcal{X})$ ,  $\mathcal{B}(\mathcal{P})$ ,  $\mathcal{B}(\mathcal{U}(x))$  and  $\mathcal{B}(\mathcal{R}_c)$ . The history random variable follows the recursion

$$H_t := (X_0, \Theta_0, U_0, R_0, \dots, X_{t-1}, \Theta_{t-1}, U_{t-1}, R_{t-1}, X_t), \quad (4)$$

<sup>5</sup> Yet, there can be alternative settings where an exogenous random variable impacts both  $R_t$  and  $X_{t+1}$ , resulting in dependence between the reward and the next state.

---

with  $(X_t, U_t) \in \mathbb{K}$ , where  $t \in \mathbb{N}_{\geq 2}$  denotes the period. The history vector's realization is denoted by

$$h_t := (x_0, \theta_0, u_0, r_0, \dots, x_{t-1}, \theta_{t-1}, u_{t-1}, r_{t-1}, x_t),$$

similarly, with  $(x_t, u_t) \in \mathbb{K}$ . In addition,  $\overline{\mathcal{H}}_0 := \mathcal{H}_0 = \mathcal{X}$ , and for  $t \geq 1$ ,  $\mathcal{H}_t$  is a subspace of

$$\overline{\mathcal{H}}_t := (\mathcal{X} \times \mathcal{P} \times \mathcal{U} \times \mathcal{R}_c)^t \times \mathcal{X} = \mathcal{H}_{t-1} \times \mathcal{P} \times \mathcal{U} \times \mathcal{R}_c \times \mathcal{X}.$$

The infinite sequence of four-tuples  $\overline{\mathcal{H}}_\infty$  is the sample space, denoted by  $\Omega$ ,

$$\Omega = \overline{\mathcal{H}}_\infty := (\mathcal{X} \times \mathcal{P} \times \mathcal{U} \times \mathcal{R}_c)^\infty = \mathcal{X} \times \mathcal{P} \times \mathcal{U} \times \mathcal{R}_c \times \mathcal{X} \times \mathcal{P} \times \mathcal{U} \times \mathcal{R}_c \dots$$

$\Omega$  is the space of histories  $\overline{H}_t = (X_0, \Theta_0, U_0, R_0, X_1, \Theta_1, U_1, R_1, \dots)$  with  $X_t \in \mathcal{X}$ ,  $\Theta_t \in \mathcal{P}$ ,  $U_t \in \mathcal{U}$ ,  $R_t \in \mathcal{R}_c$  for  $t \geq 0$ . The state, parameter, control and reward variables are defined as projections from  $\Omega$  to sets  $\mathcal{X}$ ,  $\mathcal{P}$ ,  $\mathcal{U}$  and  $\mathcal{R}_c$ , respectively.

A typical element of the sample space,  $\omega \in \Omega$ , is an infinite sequence of the form below:

$$\omega = (x_0, \theta_0, u_0, r_0, x_1, \theta_1, u_1, r_1, \dots) \text{ where } x_t \in \mathcal{X}, \theta_t \in \mathcal{P}, u_t \in \mathcal{U}, r_t \in \mathcal{R}_c, \quad \forall t \geq 0.$$

**Definition.** The posterior distribution  $(\pi_t(\cdot | H_t) : \mathcal{P} | \mathcal{H}_t \rightarrow \mathbb{R}_+)$  is a belief distribution over  $\Theta_t \in \mathcal{P}$ . It is a function of the random and time-dependent history  $H_t$ .

**Definition.** A randomized policy  $\mu = \{\mu_t\}$  is a sequence of stochastic kernels  $\mu_t$  on  $\mathcal{U}$  given  $\mathcal{H}_t$  and  $\mathcal{P}$ , satisfying

$$\mu_t(\mathcal{U}(x_t) | h_t, \theta_t) = 1 \text{ for all } h_t \in \mathcal{H}_t, \theta_t \in \mathcal{P}, \text{ and } t \geq 0.$$

where  $\mu_t$  is an element of the set of admissible control policies, denoted by  $\mathcal{M}$ .

$\mathcal{B}(\Omega)$  is the corresponding product  $\sigma$ -algebra of  $\Omega$ . The final element of the probability space is the probability measure  $\mathbb{P}_{x_0}^{\mu, \theta} : \mathcal{B}(\Omega) \rightarrow [0, 1]$ . It represents the probability measure when policy  $\mu \in \mathcal{M}$  is used, the initial state is  $X_0 = x_0$ , and the true parameter is  $\theta$ . The expectation operator with respect to  $\mathbb{P}_{x_0}^{\mu, \theta}$  is  $\mathbb{E}_{x_0}^{\mu, \theta}$ . Whenever this expectation is taken, we take the random variables  $(X_n, \Theta_n, U_n, R_n)_{\forall n}$  as generated by  $\mathbb{P}_{x_0}^{\mu, \theta}$ . We have a collection of sample paths that are parametrized by  $\mu$ ,  $\theta$  and  $x_0$ . The operands of these operators are specified by the underlying state process induced by  $\mu$ ,  $\theta$  and  $x_0$ . We emphasize that the space of admissible histories,  $\mathcal{H}_t$ , is contained in  $\Omega = \overline{\mathcal{H}}_\infty$ , and therefore, the (admissible) history random variable  $H_t$  is defined on  $(\Omega, \mathcal{B}(\Omega), \mathbb{P}_{x_0}^{\mu, \theta})$ .

---

A randomized, history-dependent policy  $\mu$  induces a probability measure  $\mathbb{P}_{x_0}^{\mu, \theta}$  on  $(\Omega, \mathcal{B}(\Omega))$ . By the Ionescu-Tulcea Theorem, proved in Proposition 7.28 of [Shreve \(1978\)](#), for any given policy  $\mu = \{\mu_t\} \in \mathcal{M}$ , any initial state  $X_0 = x_0$  and true parameter  $\theta \in \mathcal{P}$ , there exists a unique probability measure  $\mathbb{P}_{x_0}^{\mu, \theta}$  on  $(\Omega, \mathcal{B}(\Omega))$ , satisfying:

- (a)  $\mathbb{P}_{x_0}^{\mu, \theta}(\mathcal{H}_\infty) = 1$ ,
- (b)  $\mathbb{P}_{x_0}^{\mu, \theta}(X_0 = x_0) = 1$ ,
- (c)  $\mathbb{P}_{x_0}^{\mu, \theta}(\Theta_t = \theta_t \mid h_t) = \pi_t(\theta_t \mid h_t)$  for all  $\theta_t \in \mathcal{P}$  given  $h_t \in \mathcal{H}_t$  and  $t \geq 0$ ,
- (d)  $\mathbb{P}_{x_0}^{\mu, \theta}(U_t \in \mathbf{U} \mid h_t, \theta_t) = \mu_t(\mathbf{U} \mid h_t, \theta_t)$  for all  $\mathbf{U} \in \mathcal{B}(\mathcal{U})$  given  $h_t \in \mathcal{H}_t$ ,  $\theta_t \in \mathcal{P}$ , and  $t \geq 0$ ,
- (e)  $\mathbb{P}_{x_0}^{\mu, \theta}(R_t \in \mathbf{R} \mid h_t, \theta_t, u_t) = \int_{\mathbf{R}} f^\theta(r \mid x_t, u_t) d\lambda(r)$  for all  $\mathbf{R} \in \mathcal{B}(\mathcal{R}_c)$  given  $h_t \in \mathcal{H}_t$ ,  $\theta_t \in \mathcal{P}$ ,  $u_t \in \mathcal{U}(x_t)$ , and  $t \geq 0$ . When conditioned on  $x_t$  and  $u_t$ ,  $R_t$  is independent of  $\theta_t$  and  $h_t \setminus \{x_t\}$ .
- (f)  $\mathbb{P}_{x_0}^{\mu, \theta}(X_{t+1} \in \mathbf{X} \mid h_t, \theta_t, u_t, r_t) = \int_{\mathbf{X}} q^\theta(y \mid x_t, u_t) d\eta(y)$  for all  $\mathbf{X} \in \mathcal{B}(\mathcal{X})$  given  $h_t \in \mathcal{H}_t$ ,  $\theta_t \in \mathcal{P}$ ,  $u_t \in \mathcal{U}(x_t)$ ,  $r_t \in \mathcal{R}_c$ , and  $t \geq 0$ . When conditioned on  $x_t$  and  $u_t$ ,  $X_{t+1}$  is independent of  $\theta_t$ ,  $r_t$  and  $h_t \setminus \{x_t\}$ .

The probability measure  $\mathbb{P}_{x_0}^{\mu, \theta}$  induced by the policy  $\mu$  satisfies all (a)-(f), where

- (a) is by the definition of probability measure.
- (b) is by construction, it implies the initial state of the process is  $x_0$  with probability 1.
- (c) shows the history-dependent posterior distribution, from which the sample  $\theta_t$  is generated.
- (d) is the decision rule, i.e., the collection of policies. In each period  $t$ , the DM selects controls not only by the current state  $x_t$ , but by the entire history vector  $h_t$ . The decision rule also depends on the sample  $\theta_t$ . Since  $h_t$  ends with  $x_t$ , we add  $\theta_t$  as a condition.
- (e) characterizes the random reward drawn from the distribution which knows the noisy version of  $\theta$ . Given  $x_t$  and  $u_t$ , the random reward  $R_t$ , generated from density  $f^\theta(\cdot \mid x_t, u_t)$ , does not depend on the sample  $\theta_t$ . Since  $\theta_t$  does not give any additional information, it can be dropped. If not conditioned on  $x_t$  and  $u_t$ , then  $R_t$  depends on  $h_t$ ,  $\theta_t$ , and  $u_t$ .
- (f) is the state transition law. Set  $\mathbf{X}$  represents the states that are accessible from  $x_t$ . Given  $x_t$  and  $u_t$ , the random next state  $X_{t+1}$ , generated from density  $q^\theta(\cdot \mid x_t, u_t)$ , is independent of  $\theta_t$  and  $r_t$ . Since  $\theta_t$  and  $r_t$  are superfluous, they can be dropped. If not conditioned on  $x_t$  and  $u_t$ , then  $X_{t+1}$  depends on  $h_t$ ,  $\theta_t$ ,  $u_t$  and  $r_t$ .

---

When the expectation is over one step instead of the entire process, we use a different notations, i.e., different than  $\mathbb{E}_{x_0}^{\mu, \theta}$ . If the expectation is taken with respect to the random reward, we denote it by  $E_{f^\theta}[\cdot | x, u]$ , where density  $f^\theta$  is with respect to  $\sigma$ -finite measure  $\lambda$ . If the expectation is with respect to the random next state, the operator is  $E_{q^\theta}[\cdot | x, u]$ , where density  $q^\theta$  is with respect to the  $\sigma$ -finite measure  $\eta$ . If it is with respect to both the random reward and the random next state, then we use  $E_{f^\theta q^\theta}[\cdot | x, u]$ . Whenever there is a  $\cdot$  in these expectation operators, the  $\cdot$  implies a random variable.

### 3.2.3 Objective Function (Performance Criteria).

Given the initial state  $x_0$  and the discount factor  $\beta \in (0, 1)$ , the expected discounted reward over the infinite-horizon of implementing a policy  $\mu$  from period  $t = 0$  onward is

$$V_{x_0}^{\mu, \theta}(0) := \mathbb{E}_{x_0}^{\mu, \theta} \left[ \sum_{t=0}^{\infty} \beta^t R_t \right], \quad \forall \mu \in \mathcal{M}, x_0 \in \mathcal{X}. \quad (5)$$

When the rewards from period 0 up until  $n - 1$  are dismissed and the discounting starts from period  $n$  onward, we have

$$V_{x_0}^{\mu, \theta}(n) := \mathbb{E}_{x_0}^{\mu, \theta} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t \right]. \quad (6)$$

As in (5), the DM implements policy  $\mu$ , starting from (known) state  $x_0$  in period 0. However, the rewards in (6) are accumulated only from period  $t = n$  onward, when the (random) system state is  $X_n$ . We will use the notation  $n$  when we fix a specific time period and  $t$  when we take summations.

For a given policy, the DM should know the value of  $\theta$  to compute (5) and (6). Alternatively, the DM can statistically estimate these quantities if they have access to the reward density and transition density of the oracle to simulate the policy.

Recall from (3) that the optimal value function of the standard MDP problem is  $\nu^\theta(x)$ , with optimal policy  $\mu^\theta$ . In the adaptive setting, when the DM is assumed to know  $\theta$ , the optimal policy is  $\mu^\theta$ . Hence, without loss of generality, we define  $\nu^\theta(x_0)$  in (7) with respect to  $\mathbb{E}_{x_0}^{\mu^\theta}$ , i.e.,

$$\nu^\theta(x_0) := \sup_{\mu \in \mathcal{M}} V_{x_0}^{\mu, \theta}(0) = \sup_{\mu \in \mathcal{M}} \mathbb{E}_{x_0}^{\mu, \theta} \left[ \sum_{t=0}^{\infty} \beta^t R_t \right] = \mathbb{E}_{x_0}^{\mu^\theta, \theta} \left[ \sum_{t=0}^{\infty} \beta^t R_t \right], \quad \forall x_0 \in \mathcal{X}, \quad (7)$$

where  $\nu^\theta(x_0)$  is equal to  $\nu^\theta(x)$  in (3) when  $x = x_0, \forall x \in \mathcal{X}$ .

The process starts from state  $x_0$  in period 0. We call  $\mu^\theta$  the  $\theta$ -optimal policy. Under Assumption 0, this policy is stationary, deterministic, and Markovian.

---

## 4 Thompson Sampling

Given the formulation of the adaptive control problem, there is flexibility in how the evolution of the stochastic process can be “customized”. The specification arises two-fold: by the posterior update and the control selection. Until now, we denoted a general admissible policy by  $\mu = \{\mu_t\}$  and the belief update rule by  $\pi_t(\cdot | h_t)$ . From here onward, we adopt a particular policy, Thompson sampling (TS). We define the TS policy by specifying the decision rule  $\mu$  and the function  $\pi_t(\cdot | h_t)$  to update the belief vector. While the DM chooses  $\mu$ , the system designer determines  $\pi_t$ . We use TS, TS algorithm, TS decision rule, and TS policy interchangeably throughout the paper. Also, we denote TS by  $\tau$ , which we will formally define later in this section. The probability measure induced by TS and its corresponding expectation operator are denoted by  $\mathbb{P}_{x_0}^{\tau, \theta}$  and  $\mathbb{E}_{x_0}^{\tau, \theta}$ , respectively.

### 4.1 Posterior Update

The TS algorithm generates an estimate  $\theta_t$  in each period, by using the “synthetic” belief update function  $\pi_t(\theta_t | H_t)$ . Initially, the DM holds the prior belief  $\pi_0(\theta_0 | h_0) > 0$  on the true parameter  $\theta$ . That is, the unknown  $\theta$  is modeled by the  $|\mathcal{P}|$ -valued random variable  $\Theta_t$ , with initial prior distribution

$$\pi_0(\theta_0 | h_0) := \mathbb{P}_{x_0}^{\tau, \theta}(\Theta_0 = \theta_0 | h_0), \quad \forall \theta_0 \in \mathcal{P}.$$

At the beginning of period  $t$ , the DM updates her belief over the parameter candidates, by computing the (random) posterior distribution

$$\pi_t(\theta_t | H_t) := \mathbb{P}_{x_0}^{\tau, \theta}(\Theta_t = \theta_t | H_t), \quad \forall \theta_t \in \mathcal{P}. \quad (8)$$

The expected value of  $\pi_t(\theta_t | H_t)$ ,

$$\pi_t(\theta_t) := \mathbb{E}_{x_0}^{\tau, \theta}[\pi_t(\theta_t | H_t)],$$

is deterministic, and its dependence on  $(x_0, \tau, \theta)$  is implicit. We employ Bayes’ Theorem to conduct the update,

$$\pi_t(\theta_t | H_t) := \frac{\mathcal{L}_t^{\theta_t}(H_t)\pi_0(\theta_0 | h_0)}{\sum_{\gamma \in \mathcal{P}} \mathcal{L}_t^\gamma(H_t)\pi_0(\gamma | h_0)}, \quad (9)$$

where  $\mathcal{L}_t^\gamma(H_t) : \mathcal{H}_t \rightarrow \mathbb{R}$  is the (history-dependent) likelihood function. For any  $\gamma \in \mathcal{P}$ ,

$$\mathcal{L}_t^\gamma(H_t) := \prod_{s=1}^t f^\gamma(R_{s-1} | X_{s-1}, U_{s-1})q^\gamma(X_s | X_{s-1}, U_{s-1}).$$

The joint density  $f^\gamma(\cdot | x, u)q^\gamma(\cdot | x, u)$  specifies a joint probability measure on  $[0, 1] \times \mathcal{X}$ ,

$$\rho_{x,u}^\gamma(\mathbf{R}, \mathbf{X}) := \int_{\mathbf{R}} f^\gamma(r | x, u) d\lambda(r) \int_{\mathbf{X}} q^\gamma(y | x, u) d\eta(y),$$

for  $\mathbf{R} \subseteq \mathcal{B}(\mathcal{R}_c)$ ,  $\mathbf{X} \subseteq \mathcal{B}(\mathcal{X})$ . Then, for any parameter value  $\gamma \in \mathcal{P}$ , the ratio of the Radon-Nikodym derivative is

$$\frac{d\rho_{x,u}^\theta}{d\rho_{x,u}^\gamma} = \frac{f^\theta(\cdot | x, u)q^\theta(\cdot | x, u)}{f^\gamma(\cdot | x, u)q^\gamma(\cdot | x, u)}.$$

**Definition.** The relative entropy of  $\rho_{x,u}^\theta$  with respect to  $\rho_{x,u}^\gamma$  is

$$\mathcal{K}(\rho_{x,u}^\theta | \rho_{x,u}^\gamma) := E_{f^\theta q^\theta} \left[ \log \left( \frac{d\rho_{x,u}^\theta}{d\rho_{x,u}^\gamma} \right) \right],$$

given  $\rho_{x,u}^\theta$  is absolutely continuous with respect to  $\rho_{x,u}^\gamma$ .

Note that the expectation operator  $E_{f^\theta q^\theta}[\cdot | x, u]$  was defined earlier in Section 3.2, such that we integrate over the random reward and next state, for one step only. Next, we specify the decision rule to characterize the TS policy.

## 4.2 Decision Rule

**Definition.** The Thompson sampling policy  $\tau = \{\tau_t\}$  is a sequence of stochastic kernels  $\tau_t$  on  $\mathcal{U}$  given  $h_t$  and  $\theta_t$ , satisfying

$$\tau_t(\cdot | h_t, \theta_t) := \mu^{\theta_t}(\cdot | x_t).$$

Under Assumption 0,  $\mu^{\theta_t}$  is a stationary, deterministic, and Markovian policy. Since  $\mu^{\theta_t}$  only depends on  $x_t$ , (by definition) the  $\tau$  policy depends on  $x_t$ , instead of  $h_t$ . In each period  $t$ , the TS decision rule samples  $\theta_t$  and employs  $\mu^{\theta_t}$ , i.e., it picks the control that maximizes the expected infinite-horizon discounted reward by treating  $\theta_t$  as the true value of the unknown parameter  $\theta$ . The TS decision rule is deterministic, given  $\theta_t$ . However, until the (random and history-dependent) sample  $\Theta_t$  is drawn from the posterior distribution  $\pi_t(\theta_t | H_t)$ , it is a randomized decision rule.

Recall the evolution of the stochastic process from Section 3.2.1. Based on the state-control pair, the DM observes a noisy reward generated by  $f^\theta(\cdot | x_t, u_t)$ , and thus, cannot immediately identify the true value  $\theta$ . Then, through the transition density  $q^\theta(\cdot | x_t, u_t)$ , the current state transitions into the next state. After observing the reward and the transition, the DM updates the posterior on every  $\gamma \in \mathcal{P}$  using (9). Afterwards, a new parameter estimate  $\theta_{t+1}$  is sampled from the updated distribution, leading to the next control  $u_{t+1}$ .

---

We underline that the history vector includes the sample, i.e.,  $H_t := (X_0, \Theta_0, U_0, R_0, \dots)$  contains the period- $t$  sample  $\theta_t$ , which renders the TS policy well defined on the probability space  $(\Omega, \mathcal{B}(\Omega), \mathbb{P}_{x_0}^{\tau, \theta})$ , defined in Section 3.2.

**Lemma 4.1** (Degenerate prior). *TS is equivalent to the  $\theta$ -optimal policy when the prior distribution is degenerate, i.e.,  $\pi_0(\theta | h_0) = 1$ .*

We defer the proof of Lemma 4.1 to Appendix A.

## 5 Decomposition of Expected Regret

The objective of this section is to decompose the standard regret into interpretable components and illustrate how the third component, i.e., residual regret, decays to 0 through a numerical example. Because we are in an infinite-horizon setting, we can consider expected regret in any given fixed period  $n$  in the future, as assessed in period 0. In what follows, we decompose this “expected regret process” into three different components, two of which that cannot be changed from period  $n$  onward and a third which can be influenced. The first component is the traditional notion of expected finite-horizon regret.

### 5.1 Standard Notion of Expected Regret

One would naturally quantify the expected regret of an admissible policy by taking the difference between the optimal value function and the TS policy’s value function. By the definitions in (5) and (7), this difference is equal to

$$\begin{aligned} E[\text{Regret}^\theta(0, \infty)] &:= \nu^\theta(x_0) - V_{x_0}^{\tau, \theta}(0) \\ &= \mathbb{E}_{x_0}^{\mu^\theta, \theta} \left[ \sum_{t=0}^{\infty} \beta^t R_t \right] - \mathbb{E}_{x_0}^{\tau, \theta} \left[ \sum_{t=0}^{\infty} \beta^t R_t \right], \end{aligned} \quad (10)$$

where  $(0, \infty)$  represents the starting and ending periods, inclusive. The expected infinite-horizon regret  $E[\text{Regret}^\theta(0, \infty)]$  is a function of two different expectation operators, so the expectation “ $E$ ” represents a label rather than a formal mathematical expression. We emphasize that even though  $R_t$  appears in both terms of (10), one of them is driven by the process generated by  $\mu^\theta$ , while the other one is driven by  $\tau$ .

By construction,  $E[\text{Regret}^\theta(0, \infty)]$  can be partitioned into two components; a finite component

that tallies rewards up until some period  $n - 1$ , and the remainder that goes into infinity, i.e.,

$$E[\text{Regret}^\theta(0, \infty)] := E[\text{Regret}^\theta(0, n - 1)] + E[\text{Regret}^\theta(n, \infty)]. \quad (11)$$

When  $\beta \in [0, 1)$ , the difference in (10) is the expected value of the regret felt by the DM discounted back to period 0, i.e., in period-0 “dollars”. To express the regret of the  $\theta$ -MDP in period- $n$  dollars, we multiply (10) with  $\beta^{-n}$ , to obtain

$$E[\text{Regret}_n^\theta(0, \infty)] := (\nu^\theta(x_0) - V_{x_0}^{\tau, \theta}(0))\beta^{-n} \quad (12)$$

$$= \left[ \mathbb{E}_{x_0}^{\mu^\theta, \theta} \left[ \sum_{t=0}^{\infty} \beta^t R_t \right] - \mathbb{E}_{x_0}^{\tau, \theta} \left[ \sum_{t=0}^{\infty} \beta^t R_t \right] \right] \beta^{-n} \quad (13)$$

$$= \left[ \mathbb{E}_{x_0}^{\mu^\theta, \theta} \left[ \sum_{t=0}^{n-1} \beta^t R_t \right] - \mathbb{E}_{x_0}^{\tau, \theta} \left[ \sum_{t=0}^{n-1} \beta^t R_t \right] \right] \beta^{-n} \quad (14)$$

$$+ \left[ \mathbb{E}_{x_0}^{\mu^\theta, \theta} \left[ \sum_{t=n}^{\infty} \beta^t R_t \right] - \mathbb{E}_{x_0}^{\tau, \theta} \left[ \sum_{t=n}^{\infty} \beta^t R_t \right] \right] \beta^{-n}. \quad (15)$$

The subscript  $n$  in  $E[\text{Regret}_n^\theta(0, \infty)]$  corresponds to the period at which the money is evaluated. This shows that the decomposition (11) also applies to the regret expressed in any period- $n$  dollars.

$$E[\text{Regret}_n^\theta(0, \infty)] := E[\text{Regret}_n^\theta(0, n - 1)] + E[\text{Regret}_n^\theta(n, \infty)].$$

When  $\beta = 1$ , there is no difference between (10) and (13). We decompose (13) into two components, namely (14) and (15). In particular, (14) represents the expected finite-horizon regret, and by dividing it by  $\beta^n$  we convert it to period- $n$  dollars. We denote it by  $E[\text{Regret}_n^\theta(0, n - 1)]$ . Recall that  $\mu^\theta$  is an optimal policy for the infinite-horizon discounted reward maximization problem, with value  $\nu^\theta(x_0)$ . When  $\beta = 1$ ,  $E[\text{Regret}_n^\theta(0, n - 1)]$  becomes the traditional expected regret over a finite horizon. We observe from Example 2.1 that the non-discounted version of  $E[\text{Regret}_n^\theta(0, n - 1)]$  violates the order of  $\ln(n)$ . The expected regret component (15) can be interpreted as the expected infinite-horizon regret, starting in period  $n$ , in period- $n$  dollars. Distributing  $\beta^{-n}$  inside, (15) can alternatively be written as

$$\mathbb{E}_{x_0}^{\mu^\theta, \theta} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t \right] - \mathbb{E}_{x_0}^{\tau, \theta} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t \right]. \quad (16)$$

By the first remark in Section 3.1, the infinite geometric series property, and assuming  $\beta \in [0, 1)$ , (16) is upper bounded by  $\frac{2M}{1-\beta}$ , which is a constant independent of  $n$ . When  $\beta = 1$ , (16) may grow to infinity in  $n$ .

We revisit Example 2.1, which was introduced in Section 2.2. The value of the  $\tau$  policy  $V_{x_0}^{\tau, \theta}(0)$  is equal to the probability that the initial sample is equal to the true parameter  $\theta$  times the infinite-horizon reward of having the first guess right. In the case of sampling the true parameter at  $t = 0$ , the DM earns a reward of 1 in all periods, except for the first period ( $t = 0$ ). In this case, the total discounted reward accrued starting at  $t = 1$  is  $\frac{1}{1-\beta}$ . By the definition of  $V_{x_0}^{\tau, \theta}(0)$ , we convert the total reward into period-0 dollars by multiplying  $\frac{1}{1-\beta}$  with  $\beta$ . Therefore,

$$V_{x_0}^{\tau, \theta}(0) = \pi_0(\theta | h_0) \frac{\beta}{1 - \beta}.$$

The value of the  $\theta$ -optimal policy in period-0 dollars is

$$\nu^\theta(x_0) = \frac{\beta}{1 - \beta}.$$

Hence, by (13) the expected infinite-horizon regret of the  $\tau$  policy is

$$E[\text{Regret}_n^\theta(0, \infty)] := (\nu^\theta(x_0) - V_{x_0}^{\tau, \theta}(0))\beta^{-n} = \left( (1 - \pi_0(\theta | h_0)) \frac{\beta}{1 - \beta} \right) \beta^{-n}.$$

We now inspect the limiting behavior of this metric (as  $n \rightarrow \infty$ ). When the discount factor  $0 \leq \beta < 1$ , we have

$$\lim_{n \rightarrow \infty} E[\text{Regret}_n^\theta(0, \infty)] = \lim_{n \rightarrow \infty} \left( (1 - \pi_0(\theta | h_0)) \frac{\beta}{1 - \beta} \right) \beta^{-n} = \infty. \quad (17)$$

The expected finite-horizon regret (14) of the  $\tau$  policy is

$$\begin{aligned} E[\text{Regret}_n^\theta(0, n-1)] &:= \left( \mathbb{E}_{x_0}^{\mu^\theta, \theta} \left[ \sum_{t=0}^{n-1} \beta^t R_t \right] - \mathbb{E}_{x_0}^{\tau, \theta} \left[ \sum_{t=0}^{n-1} \beta^t R_t \right] \right) \beta^{-n} \\ &= \left( \sum_{t=1}^{n-1} \beta^t - \pi_0(\theta | h_0) \sum_{t=1}^{n-1} \beta^t \right) \beta^{-n} \\ &= \left( \sum_{t=1}^{n-1} \beta^t (1 - \pi_0(\theta | h_0)) \right) \beta^{-n} \\ &= \frac{\beta(1 - \beta^{n-1})\beta^{-n}}{1 - \beta} (1 - \pi_0(\theta | h_0)) = \frac{\beta^{1-n} - 1}{1 - \beta} (1 - \pi_0(\theta | h_0)). \end{aligned} \quad (18)$$

Similar to (17), we inspect the limiting behavior of the expected finite-time regret, which yields

$$\lim_{n \rightarrow \infty} E[\text{Regret}_n^\theta(0, n-1)] = \lim_{n \rightarrow \infty} \frac{\beta^{1-n} - 1}{1 - \beta} (1 - \pi_0(\theta | h_0)) = \infty.$$

When  $\beta = 1$  (18) is equal to  $(n-1)(1 - \pi_0(\theta | h_0))$ , thus the expected finite-time regret grows linearly, not logarithmically as in Agrawal and Goyal (2012).

---

## 5.2 Components of Expected Regret

We now propose an alternative decomposition of  $E[\text{Regret}_n^\theta(0, \infty)]$ . By construction, the second term in (16) is equal to the value function of the  $\tau$  policy, i.e.,  $V_{x_0}^{\tau, \theta}(n)$ . Moreover, the first term in (16) is equal to the expectation of the optimal value function with respect to the  $\theta$ -optimal policy, i.e.,

$$\mathbb{E}_{x_0}^{\mu^\theta, \theta}[\nu^\theta(X_n)] := \mathbb{E}_{x_0}^{\mu^\theta, \theta} \left[ \mathbb{E}_{X_n}^{\mu^\theta, \theta} \left[ \sum_{t=0}^{\infty} \beta^t R'_t \right] \right] = \mathbb{E}_{x_0}^{\mu^\theta, \theta} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t \right]. \quad (19)$$

In (19),  $X_n$  is generated by running the  $\mu^\theta$  policy starting in period  $t = 0$  from state  $x_0$ . Conditional on the random “starting” state  $X_n$ ,  $\{R'_0, R'_1, \dots\}$  is the random reward process generated by the optimal policy  $\mu^\theta$ . When unconditioned on  $X_n$ ,  $R'_t \sim R_{t+n}$  given starting state  $x_0$ . Adding and subtracting  $\mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)]$ , and regrouping yields the following decomposition.

$$\begin{aligned} E[\text{Regret}_n^\theta(0, \infty)] & \\ & := (\nu^\theta(x_0) - V_{x_0}^{\tau, \theta}(0))\beta^{-n} \\ & = E[\text{Regret}_n^\theta(0, n-1)] \qquad \qquad \qquad \text{(Expected finite-time regret)} \end{aligned} \quad (20)$$

$$+ \mathbb{E}_{x_0}^{\mu^\theta, \theta}[\nu^\theta(X_n)] - \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)] \qquad \qquad \qquad \text{(Expected state regret)} \quad (21)$$

$$+ \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)] - V_{x_0}^{\tau, \theta}(n) \qquad \qquad \qquad \text{(Expected residual regret).} \quad (22)$$

Notice that (20) is the same quantity as (14). Next, we discuss (21) and (22).

### 5.2.1 Expected State Regret.

The expected state regret (21) captures the unavoidable future consequences of landing in a suboptimal state after implementing TS for  $n$  periods, which we formally define below.

**Definition.** The expected state regret,

$$\mathcal{S}_{x_0}^{\tau, \theta}(n) := \mathbb{E}_{x_0}^{\mu^\theta, \theta}[\nu^\theta(X_n)] - \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)],$$

is the expected forward-looking regret under an optimal policy from period  $n$  onward. It quantifies the difference between landing in a random state  $X_n$  when the  $\tau$  policy is started in period 0 versus a potentially different random state  $X_n$ , had an optimal policy been followed instead from the starting period.

Although it tallies the difference in rewards starting from period  $n$  into the infinite future, we only denote the starting period  $n$  inside the parenthesis.




Figure 3: Illustration of two different sample paths giving rise to the expected state regret.

The random state  $X_n$  in  $\mathbb{E}_{x_0}^{\mu^\theta, \theta}[\nu^\theta(X_n)]$  is induced by running the  $\mu^\theta$  policy for  $n$  periods, whereas the random  $X_n$  in  $\mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)]$  arises under the  $\tau$  policy. Starting from the respective random states  $X_n$ , both sample paths follow the  $\theta$ -optimal policy. Figure 3 illustrates two representative sample paths. If there is no state process or if the state process is iid, then these two terms would be equal. Since our problem involves a nontrivial state process, the DM who implements the  $\tau$  policy may end up in a “bad” part of the state space, leading to an unavoidable, positive penalty. Recall that  $\nu^\theta(\cdot)$  is the optimal value function of the DM who *knows* the true  $\theta$ . After period  $n$ , the DM cannot do any better than  $\mu^\theta$ .

**Example 5.1** (Absorption into an unfavorable set of states). Consider Figure 4. Similar to Example 2.1, in period  $n = 0$ , the DM can choose either  $A$  or  $B$  with an immediate reward of 0. After being absorbed into one of  $x_A$  or  $x_B$ , the DM can pick either control 1 or 2 and receives a deterministic reward, as a function of the control and true parameter. Suppose the true parameter  $\theta$  is  $B$ , which is unknown to the DM who performs TS.

If  $\pi(A | x_0) \approx 1$ , then the DM would initially sample  $A$ , and consequently would stay in  $x_A$  forever. The  $\tau$  policy may then pick control 1 (generating a reward of 0) or pick control 2




Figure 4: Deterministic reward depending on the control, where true parameter is B.

(generating a reward of 0.5). Recall that the system designer provides the transition and reward densities, i.e., the reward structures  $R^A(\cdot)$  and  $R^B(\cdot)$  are both known to the DM. Since the rewards are deterministic and have different values, regardless of the control picked at  $n = 1$ , the  $\tau$  policy will immediately learn that  $\theta = B$ . Hence,  $\tau$  will always pick control 2 from  $n = 2$  onward, incurring a reward of 1 forever. The  $\theta$ -optimal policy will pick parameter  $B$  in period  $n = 0$ , receiving a reward of 0.5. Then, by (21), the undiscounted expected total regret of being in state  $x_A$ , as opposed to  $x_B$ , increases linearly, by  $1 - 0.5 = 0.5$  in each period. In contrast, the final term (22) equals 0 when  $n \geq 2$ . Once landing in state  $x_A$ , after 1 period both TS and an optimal policy will choose the same control, i.e., control 2, forever. However, (22) does not always converge to 0 as rapidly. In the next section, we illustrate that when the rewards are not deterministic, the  $\tau$  policy learns more slowly.

### 5.2.2 Expected Residual Regret.

Now, we will formally introduce the third component of the expected infinite-horizon expected regret, i.e., (22), and study its asymptotic behavior.

**Definition.** The expected residual regret, i.e.,  $\mathcal{R}_{x_0}^{\tau, \theta}(n)$ , is the expected forward-looking regret from period  $n$  onward into the infinite future. This regret is between a policy which implements  $\tau$  until it switches to the optimal policy  $\mu^\theta$  in period  $n$  as opposed to continuing with  $\tau$ . Formally,

$$\mathcal{R}_{x_0}^{\tau, \theta}(n) := \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)] - V_{x_0}^{\tau, \theta}(n). \quad (23)$$

Similar to the expected state regret, we only denote the starting period  $n$  inside the parenthesis.

Consider the first term of (23),

$$\mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)] := \mathbb{E}_{x_0}^{\tau, \theta} \left[ \sup_{\mu \in \mathcal{M}} V_{X_n}^{\mu, \theta}(0) \right] = \mathbb{E}_{x_0}^{\tau, \theta} \left[ \sup_{\mu \in \mathcal{M}} \mathbb{E}_{X_n}^{\mu, \theta} \left[ \sum_{t=0}^{\infty} \beta^t R_t \right] \right].$$

The  $\theta$ -optimal policy starts from a random state  $X_n$ , which is driven by running the  $\tau$  policy for  $n$  periods, starting at  $x_0$ . Recall Sections 3.2.2 and 4 for details. The expectation is taken over all paths leading to all possible  $X_n$ . Hence,  $\mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)]$  is a deterministic quantity. Consider the second term of (23),  $V_{x_0}^{\tau, \theta}(n)$ , which is equal to (6) by substituting the  $\tau$  policy,

$$V_{x_0}^{\tau, \theta}(n) := \mathbb{E}_{x_0}^{\tau, \theta} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t \right].$$

The expectation operator is induced by  $\tau$ , the starting state  $x_0$  and the true parameter  $\theta$ . Hence, similar to the first term,  $V_{x_0}^{\tau, \theta}(n)$  does not forget the past; periods 0 to  $(n-1)$  impact the state wherein the process finds itself in period  $n$ . Both  $\mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)]$  and  $V_{x_0}^{\tau, \theta}(n)$  discard the rewards generated during the first  $n$  periods; however, they are not independent of the past, since the random state  $X_n$  is driven by the tuple  $(x_0, \tau, \theta)$ . In summary, we decompose the standard regret into three components, i.e.

$$E[\text{Regret}^\theta(0, \infty)] = E[\text{Regret}_n^\theta(0, n-1)] + \mathcal{S}_{x_0}^{\tau, \theta}(n) + \mathcal{R}_{x_0}^{\tau, \theta}(n). \quad (24)$$

Notice that in (24)  $\mathcal{S}_{x_0}^{\tau, \theta}(n)$  and  $\mathcal{R}_{x_0}^{\tau, \theta}(n)$  tally the difference in rewards starting from  $n$  into infinity, while  $E[\text{Regret}^\theta(0, \infty)]$  and  $E[\text{Regret}_n^\theta(0, n-1)]$  specify both the starting and ending periods inside the parenthesis.  $E[\text{Regret}_n^\theta(0, n-1)]$  is the accumulation of the past losses, before period  $n$ . This component of the expected regret is sunk, in the sense that the DM cannot change it starting from period  $n$ . Similarly, the expected state regret  $\mathcal{S}_{x_0}^{\tau, \theta}(n)$  is the accumulation of the future losses, as a result of irrevocably being in a given state in period  $n$ , thus is also sunk.  $E[\text{Regret}_n^\theta(0, n-1)]$  and  $\mathcal{S}_{x_0}^{\tau, \theta}(n)$  measure the past and future consequences, respectively, of adopting TS from period 0 up until period  $n$ . From the perspective of a DM who is already in period  $n$ , neither can be influenced. In contrast, we regard the expected residual regret  $\mathcal{R}_{x_0}^{\tau, \theta}(n)$  as “controllable” because, starting in period  $n$ , a DM can choose a policy other than TS from that period onward. Therefore, amongst the three components, only expected residual regret represents the efficacy of continuing with  $\tau$  into the future.

Next, we illustrate how the expected residual regret quantifies the effectiveness of future decisions through an example. If the DM picks the best control(s) from period  $n$  onward, no matter how unfavorable the state  $X_n$  is, the expected residual regret starting in that period is 0.

---

### 5.2.3 Illustration of Expected Residual Regret.

To illustrate how learning occurs in a setting with stochastic rewards, consider a single-state  $\theta$ -MDP example. We show how the expected residual regret is driven down to 0 as a result of  $\tau$  learning over time.




Figure 5: Stochastic rewards depending on the control, where the true parameter is B.

**Example 5.2** (Expected residual regret converges to 0). Consider the single-state  $\theta$ -MDP in Figure 5. The underlying reward structure is illustrated on the arcs, which represent the controls. The rewards are stochastic. Based on the true parameter (which we assume is  $B$ ) and the control picked at every step, the reward is generated from a normal distribution with known mean and variance. Not knowing the current state, the DM draws a sample from the posterior distribution in each period and obtains either  $\Theta_t = A$  or  $\Theta_t = B$ . The initial prior belief on  $A$  is 0.5. In each step of the process, if the sample drawn is  $A$ , then the DM picks control 1. This is because control 1 has an expected reward of 0.5 when sampling  $A$ , while control 2 has an expected reward of 0.4. On the other hand, if the DM samples  $B$ , then they pick control 2 since the corresponding expected reward is higher.

Figure 6a illustrates the evolution of the expected belief on the “wrong” parameter  $\mathbb{E}_{x_0}^{\tau, B}[\pi_n(A)]$ , i.e., the expected posterior sampling error, averaged over 100 runs of TS policy. We observe that the DM learns the true parameter  $B$  not immediately, but after approximately 40 iterations. Beyond that point, the DM always picks control 2, in order to maximize the infinite-horizon expected total reward. Figure 6b illustrates the decline in the expected residual regret for the same example.

Consider a hybrid policy that switches from  $\tau$  to  $\mu^B$  after running  $\tau$  for  $n$  periods. Since the true parameter is  $B$ , the reward distributions follow  $N(0.3, 0.1)$  and  $N(0.8, 0.1)$  when control 1 or 2 is selected, respectively, i.e., the realizations of  $R^B(x_0, 1)$  and  $R^B(x_0, 2)$  are the rewards associated with control 1 and 2, respectively. Therefore, the expected reward of control 2 is greater than the

expected reward of control 1 and the  $\mu^B$  policy always picks control 2, i.e.,

$$\int r f^B(x_0, 2) dr = 0.8 > \int r f^B(x_0, 1) dr = 0.3.$$

From period  $n$  onward, a hybrid policy gains total expected reward of

$$\begin{aligned} \mathbb{E}_{x_0}^{\tau, B}[\nu^B(X_n)] &= \mathbb{E}_{x_0}^{\tau, B} \left[ \mathbb{E}_{X_n}^{\mu^B, B} \left[ \sum_{t=0}^{\infty} \beta^t R_t \right] \right] \\ &= \sum_{t=0}^{\infty} \beta^t 0.8 = \frac{0.8}{1-\beta} = \nu^B(x_0). \end{aligned}$$

We emphasize that  $n$  corresponds to a fixed time period, while  $t$  is used when summing over the rewards. Consider the  $\mu_1$  policy that always picks control 1. Had the DM implemented such a policy, the expected residual regret would have been

$$\begin{aligned} \mathcal{R}_{x_0}^{\mu_1, B}(n) &:= \mathbb{E}_{x_0}^{\mu_1, B}[\nu^B(X_n)] - V_{x_0}^{\mu_1, B}(n) \\ &= \frac{0.8}{1-\beta} - \mathbb{E}_{x_0}^{\mu_1, B} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t \right] \\ &= \sum_{t=0}^{\infty} \beta^t 0.8 - \sum_{t=n}^{\infty} \beta^{t-n} 0.3 = \sum_{t=0}^{\infty} \beta^t 0.5 = \frac{0.5}{1-\beta}. \end{aligned}$$

Thus,  $\frac{0.5}{1-\beta}$  is an upper bound on the expected residual regret for all  $n$ . After some number of periods, the rate at which the policy  $\tau$  picks control 1 becomes negligible. Once the policy no longer picks control 1, the DM has figured out the true parameter is  $B$ . If we call this period  $\tilde{n}$ , the expected residual regret becomes

$$\mathcal{R}_{x_0}^{\tau, B}(n) := \mathbb{E}_{x_0}^{\tau, B}[\nu^B(X_n)] - V_{x_0}^{\tau, B}(n) = \sum_{t=0}^{\infty} \beta^t 0.8 - \sum_{t=n}^{\infty} \beta^{t-n} 0.8 = 0, \quad \forall n \geq \tilde{n}.$$


Starting from  $n = 0$  and using Example 5.2 assumptions, the expected residual regret can be computed as

$$\mathcal{R}_{x_0}^{\tau, B}(n) = \sum_{t=0}^{\infty} \beta^t 0.8 - \left( \mathbb{E}_{x_0}^{\tau, B}[\pi_n(A)] \frac{0.3}{1-\beta} + \mathbb{E}_{x_0}^{\tau, B}[\pi_n(B)] \frac{0.8}{1-\beta} \right).$$


Figure 6b shows the evolution of  $\mathcal{R}_{x_0}^{\tau, B}$  for  $\beta = 0.9$ . When a different  $\beta \in [0, 1)$  is chosen, the trajectory remains the same, but the expected residual regret values are different for small  $n$ .

## 6 Analysis of Expected Residual Regret for Thompson Sampling

The goal of this section is to show that the expected residual regret of TS vanishes (i.e., converges to 0) in an exponential rate. To do so, we first relate the concept of ADO (Hernández-Lerma 2012)



(a) Evolution of expected posterior sampling error



(b) Evolution of expected residual regret

Figure 6: Evolution of expected posterior and expected residual regret in Example 5.2 when first sample is wrong

to the expected residual regret, showing that they are equivalent in our setting. This allows us to combine the machinery of Hernández-Lerma (2012) with Kim (2017) to obtain our result.

### 6.1 ADO and Expected Residual Regret

To study adaptive control problems in the discounted case, Schäl (1987) introduced an asymptotic definition of optimality, asymptotic discount optimality. Hernández-Lerma (2012) describes the idea behind it as “to allow the system to run during a learning period of  $n$  stages” and defines an asymptotically discount optimal (ADO) policy as follows:

---

**Definition (ADO).** A policy  $\mu$  is called *asymptotically discount optimal* (ADO) if,

$$\left| V_{x_0}^{\mu, \theta}(n) - \mathbb{E}_{x_0}^{\mu, \theta}[\nu^\theta(X_n)] \right| \rightarrow 0 \text{ as } n \rightarrow \infty, \quad \forall x_0 \in \mathcal{X},$$

where  $\nu^\theta(X_n)$  is defined by (7) and  $V_{x_0}^{\mu, \theta}(n)$  by (6).

We label an ADO policy as  $\theta$ -ADO to emphasize the dependence on the underlying parameter  $\theta$ . Traditionally, expected regret is formulated without the absolute value (Lai and Robbins 1985).

It follows that the expected residual regret is equivalent to the  $\theta$ -ADO expression, i.e.,

$$\mathcal{R}_{x_0}^{\tau, \theta}(n) := \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)] - V_{x_0}^{\tau, \theta}(n) = |V_{x_0}^{\tau, \theta}(n) - \mathbb{E}_{x_0}^{\mu, \theta}[\nu^\theta(X_n)]|. \quad (25)$$

Showing (25) will be instrumental in bounding the expected residual regret.

**Lemma 6.1.** *The absolute value in the  $\theta$ -ADO expression can be omitted in our setting, i.e.,*

$$|V_{x_0}^{\tau, \theta}(n) - \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)]| = \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)] - V_{x_0}^{\tau, \theta}(n).$$

The statement in Lemma 6.1 is self-evidently true, but because the notation is cumbersome in this setting, we defer the proof to Appendix A.

Rather than calling policies  $\theta$ -ADO, we say that they have “vanishing expected residual regret”.

**Definition.** A policy  $\mu$  has vanishing expected residual regret if,

$$\lim_{n \rightarrow \infty} \mathcal{R}_{x_0}^{\mu, \theta}(n) := \lim_{n \rightarrow \infty} \left[ \mathbb{E}_{x_0}^{\mu, \theta}[\nu^\theta(X_n)] - V_{x_0}^{\mu, \theta}(n) \right] = 0, \quad \forall x_0 \in \mathcal{X},$$

where  $\theta$  is the true parameter and  $X_n$  is the random period- $n$  state, which is obtained by running an admissible policy  $\mu$  for  $n$  periods.

If TS has vanishing expected residual regret, then eventually its expected performance converges to that of an optimal policy.

## 6.2 Temporal Difference Error

To bound the expected residual regret, we first establish a connection to the temporal-difference error function (Sutton and Barto 2018). This will later allow us to provide a bound for the former by bounding the latter. This function, which is parametrized by  $\theta$ , quantifies the discrepancy between the reward-to-go of choosing the optimal control instead of an arbitrary control in a given state.

---

**Definition** (Hernández-Lerma, 2012). We denote the temporal-difference error function by  $\phi^\theta : \mathbb{K} \rightarrow \mathbb{R}$ , where

$$\phi^\theta(x, u) := r^\theta(x, u) + \beta \int \nu^\theta(y) Q^\theta(dy | x, u) - \nu^\theta(x).$$

The first two terms of  $\phi^\theta(x, u)$  constitute the reward-to-go of choosing (an arbitrary) control  $u \in \mathcal{U}(x)$  in state  $x$ , while  $\nu^\theta(x)$  is the reward-to-go of choosing the optimal control in  $x$ .

**Lemma 6.2** (Temporal-difference error). *For every initial state  $x_0 \in \mathcal{X}$ , a policy  $\mu$  has vanishing expected residual regret, i.e.,*

$$\lim_{n \rightarrow \infty} \left[ \mathbb{E}_{x_0}^{\mu, \theta} [\nu^\theta(X_n)] - V_{x_0}^{\mu, \theta}(n) \right] = 0, \quad \forall x_0 \in \mathcal{X},$$

if and only if  $\phi^\theta(X_t, U_t) \rightarrow 0$  in probability- $\mathbb{P}_{x_0}^{\mu, \theta}$  for every  $x_0 \in \mathcal{X}$ .

We defer the proof of Lemma 6.2 to Appendix A, which has been adapted from Hernández-Lerma (2012) to our setting. The proof establishes a connection between the expected residual regret and the expected value of  $\phi^\theta$ . In particular, for a policy  $\mu$ ,

$$\mathcal{R}_{x_0}^{\mu, \theta}(n) = - \sum_{t=n}^{\infty} \beta^{t-n} \mathbb{E}_{x_0}^{\mu, \theta} \phi^\theta(X_t, U_t). \quad (26)$$

### 6.3 Expected Residual Regret Bounds

In this section, we provide convergence results for TS. We will bound the expected residual regret by bounding  $\mathbb{E}_{x_0}^{\tau, \theta} [\phi^\theta(X_t, U_t) | \theta_t \neq \theta]$  and  $\mathbb{E}_{x_0}^{\tau, \theta} [\phi^\theta(X_t, U_t)]$ , respectively. The DM draws a sample in each period  $t$ , represented by the random variable  $\Theta_t$ . Recall from Section 4.1, given the true parameter  $\theta \in \mathcal{P}$ ,

$$\pi_t(\theta | H_t) = \mathbb{P}_{x_0}^{\tau, \theta}(\Theta_t = \theta | H_t) \quad (27)$$

is the probability that the sample  $\Theta_t$  is equal to  $\theta$ , conditional on  $H_t$ . Since the condition is not a known  $h_t$ , (27) is also a random variable. Similarly,

$$1 - \pi_t(\theta | H_t) = \mathbb{P}_{x_0}^{\tau, \theta}(\Theta_t \neq \theta | H_t) \quad (28)$$

is the (random) probability that the sample  $\Theta_t$  is *not* equal to  $\theta$ , given (random)  $H_t$ . Taking the expectation of both sides of (28),

$$\mathbb{E}_{x_0}^{\tau, \theta} [1 - \pi_t(\theta | H_t)] = \mathbb{E}_{x_0}^{\tau, \theta} [\mathbb{P}_{x_0}^{\tau, \theta}(\Theta_t \neq \theta | H_t)], \quad (29)$$

resolves the uncertainty of  $H_t$ . By the law of iterated expectations, (29) simplifies into

$$1 - \pi_t(\theta) = \mathbb{P}_{x_0}^{\tau, \theta}(\Theta_t \neq \theta), \quad (30)$$

i.e., the posterior probability that the sample is *not* equal to the true parameter  $\theta$ . Then, by the law of total expectation, we partition the expectation of  $\phi^\theta$ , i.e.,

$$\begin{aligned} \mathbb{E}_{x_0}^{\tau, \theta}[\phi^\theta(X_t, U_t)] &= \mathbb{P}_{x_0}^{\tau, \theta}(\Theta_t \neq \theta) \mathbb{E}_{x_0}^{\tau, \theta}[\phi^\theta(X_t, U_t) \mid \Theta_t \neq \theta] \\ &\quad + \mathbb{P}_{x_0}^{\tau, \theta}(\Theta_t = \theta) \mathbb{E}_{x_0}^{\tau, \theta}[\phi^\theta(X_t, U_t) \mid \Theta_t = \theta], \end{aligned}$$

where  $U_t$  is the control that maximizes the reward-to-go by treating the sampled estimate  $\Theta_t$  as the true value of the unknown parameter  $\theta$ . We can simply rewrite the probability terms to obtain,

$$\begin{aligned} \mathbb{E}_{x_0}^{\tau, \theta}[\phi^\theta(X_t, U_t)] &= (1 - \pi_t(\theta)) \mathbb{E}_{x_0}^{\tau, \theta}[\phi^\theta(X_t, U_t) \mid \Theta_t \neq \theta] \\ &\quad + \pi_t(\theta) \mathbb{E}_{x_0}^{\tau, \theta}[\phi^\theta(X_t, U_t) \mid \Theta_t = \theta]. \end{aligned}$$

But then, when  $\Theta_t = \theta$ , by the decision rule  $\tau_t(U_t \mid h_t, \theta)$ , the optimal control (of state  $x_t$ ) is taken, and the temporal difference error  $\phi^\theta(X_t, U_t)$  becomes 0, by definition. Hence, when  $\Theta_t = \theta$ ,

$$\mathbb{E}_{x_0}^{\tau, \theta}[\phi^\theta(X_t, U_t)] = (1 - \pi_t(\theta)) \mathbb{E}_{x_0}^{\tau, \theta}[\phi^\theta(X_t, U_t) \mid \Theta_t \neq \theta]. \quad (31)$$

Note that, although we express the  $\tau$  policy as a sequence of stochastic kernels, it is a deterministic policy by Assumption 0, when conditioned on  $\theta_t$ . Before presenting the main result of this section, we first bound the expectation of  $\phi^\theta(X_t, U_t)$  in Lemma 6.3, then extend a result from Kim (2017) into our setting by Lemma 6.4.

**Lemma 6.3** (Lower bound on expected  $\phi^\theta$ ). *The expected value of  $\phi^\theta(X_t, U_t)$ , conditional on  $\Theta_t \neq \theta$ , is lower bounded by a non-positive constant, i.e.,*

$$\mathbb{E}_{x_0}^{\tau, \theta}[\phi^\theta(X_t, U_t) \mid \Theta_t \neq \theta] \geq -2M \left( \frac{1 + \beta}{1 - \beta} \right), \quad (32)$$

where  $M \geq 0$  is the upper bound on the absolute value of the expected reward, per Remark 3.1.

*Proof of Lemma 6.3.* In this proof, we slightly modify the  $U_t$  notation to indicate its dependence on the sample. Let  $U_t^{\Theta_t}$  denote the control picked when sample  $\Theta_t$  is drawn from  $\mathcal{P}$ , and let  $U_t^\theta$  represent the control picked when knowing  $\theta$ . Otherwise, we cannot distinguish the  $U_t$  controls.

$$\begin{aligned} \mathbb{E}_{x_0}^{\tau, \theta}[\phi^\theta(X_t, U_t^{\Theta_t}) \mid \Theta_t \neq \theta] &= \mathbb{E}_{x_0}^{\tau, \theta}[R_t(X_t, U_t^{\Theta_t}) + \beta \int \nu^\theta(x_{t+1}) q^\theta(dx_{t+1} \mid X_t, U_t^{\Theta_t}) d\eta \mid \Theta_t \neq \theta] \\ &\quad - \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_t) \mid \Theta_t \neq \theta], \end{aligned}$$

which is equal to

$$\begin{aligned} & \mathbb{E}_{x_0}^{\tau, \theta} \left[ R_t(X_t, U_t^{\Theta_t}) + \beta \int \nu^\theta(x_{t+1}) q^\theta(dx_{t+1} | X_t, U_t^{\Theta_t}) d\eta | \Theta_t \neq \theta \right] \\ & - \mathbb{E}_{x_0}^{\tau, \theta} \left[ R_t(X_t, U_t^\theta) + \beta \int \nu^\theta(x_{t+1}) q^\theta(dx_{t+1} | X_t, U_t^\theta) d\eta | \Theta_t \neq \theta \right]. \end{aligned}$$

We can rearrange the above expression to obtain

$$\mathbb{E}_{x_0}^{\tau, \theta} [R_t(X_t, U_t^{\Theta_t}) | \Theta_t \neq \theta] - \mathbb{E}_{x_0}^{\tau, \theta} [R_t(X_t, U_t^\theta) | \Theta_t \neq \theta] \quad (33)$$

$$+ \mathbb{E}_{x_0}^{\tau, \theta} \left[ \beta \int \nu^\theta(x_{t+1}) q^\theta(dx_{t+1} | X_t, U_t^{\Theta_t}) d\eta | \Theta_t \neq \theta \right] \quad (34)$$

$$- \mathbb{E}_{x_0}^{\tau, \theta} \left[ \beta \int \nu^\theta(x_{t+1}) q^\theta(dx_{t+1} | X_t, U_t^\theta) d\eta | \Theta_t \neq \theta \right]. \quad (35)$$

Then, we introduce the function  $g^\theta : \mathcal{X} \times \mathcal{P} \rightarrow \mathbb{R}_+$

$$g^\theta(x, \theta) := \int \nu^\theta(y) q^\theta(dy | x, U_t^\theta) d\eta, \quad (36)$$

such that (34) is

$$\beta \mathbb{E}_{x_0}^{\tau, \theta} [g^\theta(X_t, \Theta_t) | \Theta_t \neq \theta]$$

and (35) is

$$- \beta \mathbb{E}_{x_0}^{\tau, \theta} [g^\theta(X_t, \theta) | \Theta_t \neq \theta].$$

By Remark 3.1 and the infinite series property, we have

$$|\nu^\theta(x)| \leq \frac{M}{1-\beta}, \quad \forall x \in \mathcal{X}.$$

By (36),

$$|g^\theta(x, \theta)| \leq \frac{M}{1-\beta} \int q^\theta(dy | x, U_t^\theta) d\eta \leq \frac{M}{1-\beta}.$$

Taking the conditional expectation yields the same upper and lower bounds, i.e.,

$$\left| \mathbb{E}_{x_0}^{\tau, \theta} [g^\theta(x, \theta) | \Theta_t \neq \theta] \right| \leq \frac{M}{1-\beta}.$$

We lower bound (34)

$$\beta \mathbb{E}_{x_0}^{\tau, \theta} [g^\theta(X_t, \Theta_t) | \Theta_t \neq \theta] \geq \frac{-\beta M}{1-\beta},$$

and upper bound (35)

$$\beta \mathbb{E}_{x_0}^{\tau, \theta} [g^\theta(X_t, \theta) | \Theta_t \neq \theta] \leq \frac{\beta M}{1-\beta},$$

to get a lower bound on their difference,

$$\beta \mathbb{E}_{x_0}^{\tau, \theta} [g^\theta(X_t, \Theta_t) \mid \Theta_t \neq \theta] - \beta \mathbb{E}_{x_0}^{\tau, \theta} [g^\theta(X_t, \theta) \mid \Theta_t \neq \theta] \geq \frac{-2\beta M}{1 - \beta}.$$

The difference between the first two reward terms, i.e., (33), can be no smaller than  $-2M$ , thus

$$\mathbb{E}_{x_0}^{\tau, \theta} [\phi^\theta(X_t, U_t^{\Theta_t}) \mid \Theta_t \neq \theta] \geq -2M + \frac{2\beta M}{1 - \beta} = -2M \left( \frac{1 + \beta}{1 - \beta} \right).$$

■

Lemma 4 of Kim (2017) bounds the expected probability that  $\theta$  is not selected in period  $t$ . We will now extend this result to the discounted infinite-horizon framework. However, for us to be able to generalize the finite-dimensional state and control spaces that Kim (2017) deals with into Borel (possibly infinite) spaces, we need to make the following assumption.

*Assumption 1.* We assume

$$\inf_{x \in \mathcal{X}, u \in \mathcal{U}, r \in \mathcal{R}_c} f^\gamma(r \mid x, u) > 0, \quad \forall \gamma \in \mathcal{P} \quad (37)$$

and

$$\inf_{x \in \mathcal{X}, u \in \mathcal{U}, y \in \mathcal{Y}} q^\gamma(y \mid x, u) > 0, \quad \forall \gamma \in \mathcal{P}. \quad (38)$$

For any  $x \in \mathcal{X}$ ,  $u \in \mathcal{U}$ , and any distinct parameter value,  $\gamma \neq \theta \in \mathcal{P}$ , there exists a positive constant  $\epsilon(x, u, \theta, \gamma) > 0$  such that

$$\inf_{x \in \mathcal{X}, u \in \mathcal{U}} \mathcal{K}(\rho_{x,u}^\theta \mid \rho_{x,u}^\gamma) > \epsilon(x, u, \theta, \gamma). \quad (39)$$

We underline that under finite state and control spaces, Assumption 1 simplifies into the prerequisites of Kim (2017). As Kim (2017) explains, (39) in Assumption 1 ensures that given  $\theta \neq \gamma$ , the probability measures  $\rho_{x,u}^\theta$  and  $\rho_{x,u}^\gamma$  are distinguishable as measured by the relative entropy. In Appendix B we illustrate the implications of Assumption 1 through Example 5.2. Example 2.1 violates this assumption.

We further emphasize that Assumption 1 implies that the exploration is passive; that is, the DM learns everything about the environment regardless of their actions. We acknowledge that, in contrast to active exploration, passive exploration is less efficient. Yet, without making additional assumptions on the underlying Markov structure, i.e., making the paper less general, it does not appear that we can extend the result in Kim (2017) to this broader framework.

The following lemma extends a result from Kim (2017) to our setting.

---

**Lemma 6.4** (Extension of Lemma 4 of [Kim \(2017\)](#)). *Under Assumption 1, Lemma 4 of [Kim \(2017\)](#) extends to our setting. That is to say, implementing TS and starting from any  $x_0 \in \mathcal{X}$ , there exists constants  $a_\theta, b_\theta > 0$  such that*

$$\mathbb{E}_{x_0}^{\tau, \theta} [1 - \pi_t(\theta \mid H_t)] \leq a_\theta e^{-b_\theta t}, \quad (40)$$

where  $a_\theta$  and  $b_\theta$  are defined as in [Kim \(2017\)](#).

Because Lemma 6.4 is an adaptation of Lemma 4 of [Kim \(2017\)](#) into our setting, we defer it to the Appendix. By the law of iterated expectations, (40) simplifies into

$$1 - \pi_t(\theta) \leq a_\theta e^{-b_\theta t}. \quad (41)$$

We are now ready to introduce the main result of this section.

**Proposition 6.5** (Upper bound on  $\mathcal{R}_{x_0}^{\tau, \theta}(n)$ ). *When Assumption 1 holds, the expected residual regret converges to 0 exponentially fast.*

$$\mathcal{R}_{x_0}^{\tau, \theta}(n) := \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)] - V_{x_0}^{\tau, \theta}(n) \leq \frac{2M(1 + \beta)a_\theta e^{-b_\theta n}}{(1 - \beta)^2},$$

where  $a_\theta$  and  $b_\theta$  are positive constants, defined in Lemma 4 of [Kim \(2017\)](#).

*Proof of Proposition 6.5.* Equations (31), (32) and (41) together yield

$$-\mathbb{E}_{x_0}^{\tau, \theta}[\phi^\theta(X_t, U_t)] \leq \frac{2M(1 + \beta)a_\theta e^{-b_\theta t}}{1 - \beta}. \quad (42)$$

By the proof of Lemma 6.2,  $\phi^\theta(X_t, U_t)$  and  $\mathcal{R}_{x_0}^{\tau, \theta}(n)$  are related, see (26). Then, by (42), the right-hand side of (26) is upper bounded,

$$-\sum_{t=n}^{\infty} \beta^{t-n} \mathbb{E}_x^{\tau, \theta}[\phi^\theta(X_t, U_t)] \leq \sum_{t=n}^{\infty} \beta^{t-n} \frac{2M(1 + \beta)a_\theta e^{-b_\theta t}}{1 - \beta}.$$

Since  $b_\theta \geq 0$ , we have  $e^{-b_\theta t} \leq e^{-b_\theta n}$  for all  $t \geq n$ . Thus, the above equation becomes

$$-\sum_{t=n}^{\infty} \beta^{t-n} \mathbb{E}_x^{\tau, \theta}[\phi^\theta(X_t, U_t)] \leq \sum_{t=n}^{\infty} \beta^{t-n} \frac{2M(1 + \beta)a_\theta e^{-b_\theta n}}{1 - \beta}. \quad (43)$$

By (26) and applying the infinite geometric series formula to (43), we obtain

$$\begin{aligned} -\sum_{t=n}^{\infty} \beta^{t-n} \mathbb{E}_{x_0}^{\tau, \theta}[\phi^\theta(X_t, U_t)] &= \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)] - V_{x_0}^{\tau, \theta}(n) \\ &= \mathcal{R}_{x_0}^{\tau, \theta}(n) \leq \frac{2M(1 + \beta)a_\theta e^{-b_\theta n}}{(1 - \beta)^2}, \end{aligned}$$

thus showing that the upper bound on the expected residual regret decays exponentially. ■

---

## 7 Complete Learning and Probabilistic Residual Regret

The goal of this section is to define a probabilistic version of the expected residual regret (the expected value of the probabilistic version is the expected residual regret itself, i.e.,  $\mathcal{R}_{x_0}^{\tau, \theta}(n)$ ), and show the conditions under which it converges  $\mathbb{P}_{x_0}^{\tau, \theta}$ -almost surely to 0 in this framework. We show that the posterior distribution of TS converges  $\mathbb{P}_{x_0}^{\tau, \theta}$ -almost surely to a point mass at  $\theta$ . This behavior is called *complete learning*.

### 7.1 Complete Learning

Kim (2017) has shown that the expected posterior sampling error of TS converges to 0 exponentially fast (see Lemma 6.4). Our result builds on this result by using the Dominated Convergence Theorem. Formally,

*Assumption 2* (Existence of the limit of  $\pi_t(\theta | H_t)$ ). Suppose that  $\lim_{t \rightarrow \infty} \pi_t(\theta | H_t)$  exists  $\mathbb{P}_{x_0}^{\tau, \theta}$ -a.s.

By imposing this assumption, we avoid the nonexistence of the limit due to oscillation. Under Assumption 1, the expected posterior sampling error of TS vanishes. We will show that this, i.e.,  $\lim_{t \rightarrow \infty} \mathbb{E}_{x_0}^{\tau, \theta}[\pi_t(\theta | H_t)] = 1$ , and Assumption 2 together imply

$$\lim_{t \rightarrow \infty} \pi_t(\theta | H_t) = 1, \quad \mathbb{P}_{x_0}^{\tau, \theta}\text{-almost surely.}$$

The next lemma is the final piece to show the occurrence of complete learning.

**Lemma 7.1.** *Suppose that TS does not exhibit complete learning, i.e.,  $\lim_{t \rightarrow \infty} \pi_t(\theta | H_t) < 1$ ,  $\mathbb{P}_{x_0}^{\tau, \theta}$ -a.s. Then, there exists an  $n \geq 1$  s.t.*

$$\mathbb{P}_{x_0}^{\tau, \theta} \left( \lim_{t \rightarrow \infty} \pi_t(\theta | H_t) < 1 - \frac{1}{n} \right) > 0, \quad \mathbb{P}_{x_0}^{\tau, \theta}\text{-almost surely.}$$

The proof of Lemma 7.1 is in Appendix A.

**Theorem 1** (TS learns  $\theta$  as  $t \rightarrow \infty$ ). Suppose that Assumptions 1 and 2 hold. Then,

$$\lim_{t \rightarrow \infty} \pi_t(\theta | H_t) = 1, \quad \mathbb{P}_{x_0}^{\tau, \theta}\text{-almost surely.}$$

*Proof of Theorem 1.* By Lemma 6.4, Assumption 2, and the Dominated Convergence Theorem, it follows that

$$1 = \lim_{t \rightarrow \infty} \mathbb{E}_{x_0}^{\tau, \theta}[\pi_t(\theta | H_t)] = \mathbb{E}_{x_0}^{\tau, \theta} \left[ \lim_{t \rightarrow \infty} \pi_t(\theta | H_t) \right], \quad \mathbb{P}_{x_0}^{\tau, \theta}\text{-a.s.}$$

Then,

$$\begin{aligned}
1 &= \mathbb{E}_{x_0}^{\tau, \theta} \left[ \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) \right] \\
&= \mathbb{E}_{x_0}^{\tau, \theta} \left[ \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) \mid \left\{ \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) < 1 - \frac{1}{n} \right\} \right] \mathbb{P}_{x_0}^{\tau, \theta} \left( \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) < 1 - \frac{1}{n} \right) \\
&\quad + \mathbb{E}_{x_0}^{\tau, \theta} \left[ \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) \mid \left\{ \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) \geq 1 - \frac{1}{n} \right\} \right] \mathbb{P}_{x_0}^{\tau, \theta} \left( \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) \geq 1 - \frac{1}{n} \right) \\
&\leq (1 - 1/n) \mathbb{P}_{x_0}^{\tau, \theta} \left( \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) < 1 - \frac{1}{n} \right) + 1 - \mathbb{P}_{x_0}^{\tau, \theta} \left( \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) < 1 - \frac{1}{n} \right) \\
&= 1 - \frac{1}{n} \mathbb{P}_{x_0}^{\tau, \theta} \left( \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) < 1 - \frac{1}{n} \right), \quad \mathbb{P}_{x_0}^{\tau, \theta}\text{-a.s.}
\end{aligned}$$

However, by Lemma 7.1,  $\lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) < 1$  implies

$$\begin{aligned}
1 &= \mathbb{E}_{x_0}^{\tau, \theta} \left[ \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) \right] \\
&\leq 1 - \frac{1}{n} \mathbb{P}_{x_0}^{\tau, \theta} \left( \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) < 1 - \frac{1}{n} \right) < 1, \quad \mathbb{P}_{x_0}^{\tau, \theta}\text{-almost surely,}
\end{aligned}$$

leading to contradiction. For all  $n \geq 1$ , it must be that

$$\mathbb{P}_{x_0}^{\tau, \theta} \left( \lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) < 1 - \frac{1}{n} \right) = 0, \quad \mathbb{P}_{x_0}^{\tau, \theta}\text{-almost surely.}$$

Thus, we must have  $\lim_{t \rightarrow \infty} \pi_t(\theta \mid H_t) = 1$ ,  $\mathbb{P}_{x_0}^{\tau, \theta}$ -almost surely. This completes the proof.  $\blacksquare$

## 7.2 Vanishing Probabilistic Residual Regret

We define the residual regret as the ‘‘probabilistic’’ version of the expected residual regret, which was defined in Section 5.2.2. In the remainder of this section, we use the terms probabilistic residual regret and residual regret interchangeably. The residual regret almost mimics the definition in (23); however, it is a random expectation, and is obtained by conditioning on the random history  $H_n$ , generated by running TS from period 0 to  $n$ .

**Definition.** The residual regret  $\mathbb{R}_{x_0}^{\tau, \theta}(n)$  is a random expectation that represents the forward-looking regret from period  $n$  onward into the infinite future. This regret is between a policy which implements  $\tau$  until it switches to the optimal policy  $\mu^\theta$  in period  $n$  as opposed to continuing with  $\tau$ .

Formally,

$$\mathbb{R}_{x_0}^{\tau,\theta}(n) := \nu^\theta(X_n) - \mathbb{E}_{x_0}^{\tau,\theta} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t \mid H_n \right] \quad (44)$$

$$= \sup_{\mu \in \mathcal{M}} \mathbb{E}_{x_0}^{\mu,\theta} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t \mid H_n \right] - \mathbb{E}_{x_0}^{\tau,\theta} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t \mid H_n \right] \quad (45)$$

$$= \mathbb{E}_{x_0}^{\mu^\theta,\theta} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t \mid H_n \right] - \mathbb{E}_{x_0}^{\tau,\theta} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t \mid H_n \right], \quad (46)$$

where the conditioning on the random history vector is denoted explicitly.

The residual regret is constructed in a way such that its expected value is the expected residual regret itself, i.e.,

$$\mathcal{R}_{x_0}^{\tau,\theta}(n) = \mathbb{E}_{x_0}^{\tau,\theta}[\mathbb{R}_{x_0}^{\tau,\theta}(n)]. \quad (47)$$

Recall the interpretation of the first term of (46); the stochastic process is driven by a “hybrid” policy that follows the  $\tau$  policy for the first  $n$  periods, and then adopts the optimal policy  $\mu^\theta$  in period  $n$  after the oracle reveals  $\theta$ . It is a random quantity due to the random starting state  $X_n$  in period  $n$ , induced by TS. The second term is the random expected infinite sum obtained by implementing TS, starting from random state  $X_n$ , and omitting the rewards from period 0 to  $n-1$ .

Next, we show that the residual regret converges to 0 in the limit  $\mathbb{P}_{x_0}^{\tau,\theta}$ -a.s., assuming it exists.

*Assumption 3* (Existence of the limit of  $\mathbb{R}_{x_0}^{\tau,\theta}(n)$ ). Suppose that  $\lim_{n \rightarrow \infty} \mathbb{R}_{x_0}^{\tau,\theta}(n)$  exists  $\mathbb{P}_{x_0}^{\tau,\theta}$ -a.s.

From (44) observe that the residual regret is the difference of two terms. Consider the first term  $\nu^\theta(X_n)$ . Under some technical conditions, if the limit of the state process  $\{X_n\}$  exists, then the limit of  $\nu^\theta(X_n)$  exists. However, since we do not impose any restrictions on the underlying chain structure and the underlying stochastic process is history-dependent, we cannot guarantee that  $\lim_{n \rightarrow \infty} X_n$  exists, and thus, there could be cases where  $\lim_{n \rightarrow \infty} \nu^\theta(X_n)$  does not converge<sup>6</sup>. Therefore, we directly assume that the limit of the residual regret exists. Similar to the expected residual regret, the following corollary shows that under certain conditions the probabilistic residual regret also vanishes in the limit.

**Corollary 7.2.** *When the conditions of Proposition 6.5 and Assumption 3 hold, the residual regret of TS vanishes, i.e.,*

$$\lim_{n \rightarrow \infty} \mathbb{R}_{x_0}^{\tau,\theta}(n) = 0, \quad \mathbb{P}_{x_0}^{\tau,\theta}\text{-almost surely.}$$

<sup>6</sup> We do not require aperiodicity; the chain could be periodic, leading to oscillating rewards.

*Proof of Corollary 7.2.* Recall that the probabilistic residual regret converges  $\mathbb{P}_{x_0}^{\tau, \theta}$ -a.s. to 0 if there exists a null set  $\mathbf{A} \in \mathcal{B}(\Omega)$  with  $\mathbb{P}_{x_0}^{\tau, \theta}(\mathbf{A}) = 0$  such that the statement holds if  $\omega \notin \mathbf{A}$ . Given Assumption 3, it suffices to show that  $\forall \epsilon > 0$ ,  $\nexists \mathbf{A} \in \mathcal{B}(\Omega)$  with  $\mathbb{P}_{x_0}^{\tau, \theta}(\mathbf{A}) = \epsilon > 0$  such that  $\lim_{n \rightarrow \infty} \mathbb{R}_{x_0}^{\tau, \theta}(n) \neq 0 \forall \omega \in \mathbf{A}$ . We prove this by contradiction.

Suppose that  $\exists \mathbf{A} \in \mathcal{B}(\Omega)$  with  $\mathbb{P}_{x_0}^{\tau, \theta}(\mathbf{A}) = \epsilon' > 0$  for some  $\epsilon' > 0$  such that  $\lim_{n \rightarrow \infty} \mathbb{R}_{x_0}^{\tau, \theta}(n) \neq 0 \forall \omega \in \mathbf{A}$ . Since  $\mathbb{R}_{x_0}^{\tau, \theta}(n) > 0$  by construction, this limit is strictly positive, i.e.,  $\lim_{n \rightarrow \infty} \mathbb{R}_{x_0}^{\tau, \theta}(n) > 0 \forall \omega \in \mathbf{A}$ .

Then, for any  $\delta > 0$ , we define  $\mathbf{A}_\delta$  as the largest measurable subset of  $\mathbf{A}$  such that  $\mathbf{A}_\delta \subset \{\omega \in \mathcal{B}(\Omega) : \lim_{n \rightarrow \infty} \mathbb{R}_{x_0}^{\tau, \theta}(n) > \delta\}$ . According to Assumption 3 and the supposition in the previous paragraph,  $\exists \delta'$  such that  $\mathbb{P}_{x_0}^{\tau, \theta}(\mathbf{A}_{\delta'}) = \epsilon' > 0$ . Then, we can write

$$\lim_{n \rightarrow \infty} \mathcal{R}_{x_0}^{\tau, \theta}(n) = \lim_{n \rightarrow \infty} \mathbb{E}_{x_0}^{\tau, \theta}[\mathbb{R}_{x_0}^{\tau, \theta}(n)] \quad (48)$$

$$= \lim_{n \rightarrow \infty} \left\{ \int_{\mathbf{A}_{\delta'}} \mathbb{R}_{x_0}^{\tau, \theta}(n) d\mathbb{P}_{x_0}^{\tau, \theta}(\omega) + \int_{\mathcal{B}(\Omega) \setminus \mathbf{A}_{\delta'}} \mathbb{R}_{x_0}^{\tau, \theta}(n) d\mathbb{P}_{x_0}^{\tau, \theta}(\omega) \right\} \quad (49)$$

$$= \int_{\mathbf{A}_{\delta'}} \lim_{n \rightarrow \infty} \mathbb{R}_{x_0}^{\tau, \theta}(n) d\mathbb{P}_{x_0}^{\tau, \theta}(\omega) + \int_{\mathcal{B}(\Omega) \setminus \mathbf{A}_{\delta'}} \lim_{n \rightarrow \infty} \mathbb{R}_{x_0}^{\tau, \theta}(n) d\mathbb{P}_{x_0}^{\tau, \theta}(\omega) \quad (50)$$

$$> \delta' \epsilon' > 0, \quad (51)$$

which contradicts Proposition 6.5. By construction, the first term in (49) converges to a number strictly greater than 0, and the second term converges to some non-negative number. Because  $\mathbb{R}_{x_0}^{\tau, \theta}(n)$  exists by Assumption 3 and is finite<sup>7</sup>, by the Dominated Convergence Theorem, we can express (49) as (50). But then, (51) contradicts with Proposition 6.5. This implies that our supposition cannot hold, and thus, the residual regret converges  $\mathbb{P}_{x_0}^{\tau, \theta}$ -a.s. to 0.  $\blacksquare$

## 8 Concluding Remarks and Discussion

Fixing a finite period  $n$ , we decompose the expected infinite-horizon regret into three components. Two of the components are not actionable by the DM. The first component is the expected regret of the past. The second is the regret expected to be accrued in the future due to the random state the  $\theta$ -MDP is in period  $n$ , which is a consequence of implementing a possibly suboptimal policy and cannot be revoked by the DM. The third component, which we called the expected residual regret, captures what a rational DM would consider as expected regret. It determines whether the DM

<sup>7</sup> Since the rewards are finite-valued, in any period, the difference in rewards of the first and second terms is bounded by a constant. By the infinite geometric series property, and assuming  $\beta \in [0, 1)$ ,  $\mathbb{R}_{x_0}^{\tau, \theta}(n)$  is finite.

---

will do the best moving forward, given their state in period  $n$ . Hence, we evaluate TS by only this third component, i.e., the expected residual regret.

We reiterate that our results hold for chain settings where  $q^\gamma(\cdot | x, u)$  is strictly positive  $\forall \gamma \in \mathcal{P}$ , which rules out chains with absorbing states. Yet, by construction, the expected residual regret is a viable concept independent of the underlying chain structure; it is applicable to any structure. Hence, deriving the performance of the residual regret in broader settings is a potential direction of further research.

In our setting, we show that the expected residual regret of TS decays exponentially to 0 in the worst case (Proposition 6.5). We explore the conditions under which the posterior sampling error converges to 0 with almost sure probability (Theorem 1). At the crux of these theorems are certain sufficient conditions, which we present in Assumption 2, which rule out oscillating functions that would prevent complete learning. Finally, we show that a probabilistic version of the expected residual regret vanishes (Corollary 7.2), similar to its expected-valued version. At the crux of this corollary is the sufficiency condition that we present in Assumption 3. A future direction of research is to study the implications of Assumptions 2 and 3 in specific problem contexts. In addition to conditions on the underlying chain structure, complete learning may be a necessary but not sufficient condition for Assumption 3 to hold.

To the best of our knowledge, the notion of an ADO policy from the adaptive learning literature had not been connected to TS. By leveraging connections between these settings, we offer a novel concept of learning. Under some mild assumptions, we provide guarantees of learning in both expected and probabilistic senses in settings with general state and control spaces, using the infinite-horizon discounted reward criterion. An interesting extension of the setup in this paper is to analyze the performance of the novel concept of learning (i.e., expected residual regret) under broader settings, e.g., under general chain structures.

## Appendix

### A Proofs of Lemmas

*Proof of Lemma 4.1.* When the Bayesian update is conducted with a degenerate distribution, it returns a degenerate distribution. We show this by induction. We have  $\pi_0(\theta | h_0) = 1$  and assume

$\pi_n(\theta | H_n) = 1$ .

$$\begin{aligned} \pi_{n+1}(\theta | H_{n+1}) &:= \frac{\mathcal{L}^\theta(H_{n+1})\pi_0(\theta | h_0)}{\sum_{\gamma \in \mathcal{P}} \mathcal{L}^\gamma(H_{n+1})\pi_0(\gamma | h_0)} \\ &= \frac{\mathcal{L}^\theta(H_{n+1})\pi_0(\theta | h_0)}{\mathcal{L}^\theta(H_{n+1})\pi_0(\theta | h_0) + \sum_{\gamma \neq \theta \in \mathcal{P}} \mathcal{L}^\gamma(H_{n+1})\pi_0(\gamma | h_0)} = \frac{\mathcal{L}^\theta(H_{n+1})}{\mathcal{L}^\theta(H_{n+1})} = 1. \end{aligned}$$

Since the posterior distribution is degenerate, TS always samples the true parameter  $\theta$  from the parameter space. Therefore, the DM who runs the  $\tau$  policy ends up implementing the  $\theta$ -optimal policy.  $\blacksquare$

*Proof of Lemma 6.1.* Consider the  $\theta$ -ADO statement,  $|V_{x_0}^{\mu, \theta}(n) - \mathbb{E}_{x_0}^{\mu, \theta}[\nu^\theta(X_n)]|$ . It can be rewritten as

$$\begin{aligned} &|V_{x_0}^{\mu, \theta}(n) - \mathbb{E}_{x_0}^{\mu, \theta}[\nu^\theta(X_n)]| \tag{52} \\ &= \left| \mathbb{E}_{x_0}^{\tau, \theta} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t - \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)] \right] \right| \\ &= \left| \mathbb{E}_{x_0}^{\tau, \theta} \left[ \mathbb{E}_{x_0}^{\tau, \theta} \left( \sum_{t=n}^{\infty} \beta^{t-n} R_t \mid X_n \right) - \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n) \mid X_n] \right] \right| \\ &= \left| \mathbb{E}_{x_0}^{\tau, \theta} \left[ \mathbb{E}_{x_0}^{\tau, \theta} \left( \sum_{t=n}^{\infty} \beta^{t-n} R_t \mid X_n \right) - \nu^\theta(X_n) \right] \right| \\ &= \left| \mathbb{E}_{x_0}^{\tau, \theta} \left[ \mathbb{E}_{x_0}^{\tau, \theta} \left( \sum_{t=n}^{\infty} \beta^{t-n} R_t(X_t, U_t) \mid X_n \right) - \sup_{\mu \in \mathcal{M}} \mathbb{E}_{X_n}^{\mu, \theta} \left[ \sum_{t=0}^{\infty} \beta^t R_t(X_t, U_t) \right] \right] \right|. \tag{53} \end{aligned}$$

The second equality is by the law of iterated expectations, and the fourth by substituting in the definition of the optimal value function. Evidently, the first term of (53) is upper bounded by the second term. Thus, (53) is equal to

$$\mathbb{E}_{x_0}^{\tau, \theta} \left[ \sup_{\mu \in \mathcal{M}} \mathbb{E}_{X_n}^{\mu, \theta} \left[ \sum_{t=0}^{\infty} \beta^t R_t(X_t, U_t) \right] - \mathbb{E}_{x_0}^{\tau, \theta} \left( \sum_{t=n}^{\infty} \beta^{t-n} R_t(X_t, U_t) \mid X_n \right) \right],$$

for any initial state  $x_0 \in \mathcal{X}$ . Rewriting the optimal value function in closed form, we obtain

$$\mathbb{E}_{x_0}^{\tau, \theta} \left[ \nu^\theta(X_n) - \mathbb{E}_{x_0}^{\tau, \theta} \left( \sum_{t=n}^{\infty} \beta^{t-n} R_t(X_t, U_t) \mid X_n \right) \right],$$

which is equal to

$$\mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)] - \mathbb{E}_{x_0}^{\tau, \theta} \left[ \sum_{t=n}^{\infty} \beta^{t-n} R_t(X_t, U_t) \right] = \mathbb{E}_{x_0}^{\tau, \theta}[\nu^\theta(X_n)] - V_{x_0}^{\tau, \theta}(n).$$

$\blacksquare$

*Proof of Lemma 6.2.* This proof is an adaptation of Theorem 3.6 of [Hernández-Lerma \(2012\)](#). By the definition in Section 6.1,

$$\phi^\theta(X_t, U_t) = \mathbb{E}_{x_0}^{\mu, \theta}[R_t(X_t, U_t) + \beta\nu^\theta(X_{t+1}) - \nu^\theta(X_t) \mid H_t, U_t],$$

for any initial state  $x_0 \in \mathcal{X}$ , admissible policy  $\mu \in \mathcal{M}$ , true parameter value  $\theta$ ,  $t \geq 0$ . For any  $t \geq 1$ , history in period  $t$  is  $h_t = (x_0, \theta_0, a_0, r_0, x_1, \theta_1, a_1, r_1, \dots, x_t)$ . Multiplying by  $\beta^{t-n}$  yields

$$\beta^{t-n}\phi^\theta(X_t, U_t) = \mathbb{E}_{x_0}^{\mu, \theta}[\beta^{t-n}R_t(X_t, U_t) + \beta^{t-n+1}\nu^\theta(X_{t+1}) - \beta^{t-n}\nu^\theta(X_t) \mid H_t, U_t].$$

Taking the expectation of both sides and by the law of total expectation, we obtain

$$\mathbb{E}_{x_0}^{\mu, \theta}[\beta^{t-n}\phi^\theta(X_t, U_t)] = \mathbb{E}_{x_0}^{\mu, \theta}[\beta^{t-n}R_t(X_t, U_t) + \beta^{t-n+1}\nu^\theta(X_{t+1}) - \beta^{t-n}\nu^\theta(X_t)].$$

Summing over all  $t \geq n$  gives

$$\begin{aligned} \sum_{t=n}^{\infty} \beta^{t-n} \mathbb{E}_{x_0}^{\mu, \theta}[\phi^\theta(X_t, U_t)] &= \sum_{t=n}^{\infty} \mathbb{E}_{x_0}^{\mu}[\beta^{t-n}R_t(X_t, U_t)] \\ &\quad + \mathbb{E}_{x_0}^{\mu, \theta} \left[ \sum_{t=n}^{\infty} (\beta^{t-n+1}\nu^\theta(X_{t+1}) - \beta^{t-n}\nu^\theta(X_t)) \right]. \end{aligned}$$

The above simplifies into

$$\sum_{t=n}^{\infty} \beta^{t-n} \mathbb{E}_{x_0}^{\mu, \theta}[\phi^\theta(X_t, U_t)] = V_{x_0}^{\mu, \theta}(n) - \mathbb{E}_{x_0}^{\mu, \theta}[\nu^\theta(X_n)].$$

Recalling the result of Lemma 6.1 as,

$$-\sum_{t=n}^{\infty} \beta^{t-n} \mathbb{E}_{x_0}^{\mu, \theta}[\phi^\theta(X_t, U_t)] = \mathbb{E}_{x_0}^{\mu, \theta}[\nu^\theta(X_n)] - V_{x_0}^{\mu, \theta}(n).$$

In the limit,  $\mu$  has vanishing expected residual regret (is  $\theta$ -ADO); that is, for every  $x_0 \in \mathcal{X}$ ,

$$\lim_{n \rightarrow \infty} (\mathbb{E}_{x_0}^{\mu, \theta}[\nu^\theta(X_n)] - V_{x_0}^{\mu, \theta}(n)) = 0,$$

if and only if, for every  $x_0 \in \mathcal{X}$ ,

$$\lim_{n \rightarrow \infty} -\sum_{t=n}^{\infty} \beta^{t-n} \mathbb{E}_{x_0}^{\mu, \theta}[\phi^\theta(X_t, U_t)] = 0.$$

This is equivalent to, for every  $x_0 \in \mathcal{X}$ ,

$$\lim_{t \rightarrow \infty} -\mathbb{E}_{x_0}^{\mu, \theta}[\phi^\theta(X_t, U_t)] = 0.$$

Hence, a policy  $\mu$  has vanishing expected residual regret if and only if, for every  $x_0 \in \mathcal{X}$ ,

$$\lim_{t \rightarrow \infty} \mathbb{E}_{x_0}^{\mu, \theta}[\phi^\theta(X_t, U_t)] = 0.$$

By Theorem 4.1.4 of [Chung \(2001\)](#), if  $\mathbb{E}_{x_0}^{\mu, \theta}[\phi^\theta(X_t, U_t)]$  converges to 0, then  $\phi^\theta(X_t, U_t)$  converges to 0 in probability- $\mathbb{P}_{x_0}^{\mu, \theta}$ , for every  $x_0 \in \mathcal{X}$ , proving the forward direction of [Lemma 6.2](#). It remains to show the reverse direction to complete the proof. By the same theorem of [Chung \(2001\)](#), for a uniformly bounded sequence  $\{\phi^\theta\}$ , convergence in probability and  $\mathcal{L}^p$  (in expectation) are equivalent. By the first remark in [Section 3.1](#), it follows that  $\{\phi^\theta\}$  is bounded by some finite number. Hence,  $\phi^\theta(X_t, U_t) \rightarrow 0$  in probability- $\mathbb{P}_{x_0}^{\mu, \theta}$  implies  $\mathbb{E}_{x_0}^{\mu, \theta}[\phi^\theta(X_t, U_t)] \rightarrow 0$  as  $t \rightarrow \infty$ .  $\blacksquare$

*Proof of [Lemma 6.4](#).* We walk the reader through the proof of [Kim \(2017\)](#) while using our notation. The proof initially defines the stochastic process, for any  $\gamma \neq \theta$ ,

$$Z_t^\gamma = \sum_{s=0}^t \log \Lambda_s^\gamma,$$

where, using our notation,

$$\begin{aligned} \Lambda_0^\gamma &= 1 \\ \Lambda_s^\gamma &= \frac{f^\theta(R_{s-1} | X_{s-1}, U_{s-1})q^\theta(X_s | X_{s-1}, U_{s-1})}{f^\gamma(R_{s-1} | X_{s-1}, U_{s-1})q^\gamma(X_s | X_{s-1}, U_{s-1})}, \end{aligned}$$

for  $0 < s \leq t$ . Next, [Kim \(2017\)](#) defines filtration  $(\mathcal{H}_t : t \geq 0)$  by  $\mathcal{H}_t = \sigma(H_t)$ , where  $H_t$  is by the definition in [\(4\)](#). Then, it follows that the stochastic process  $Z_t^\gamma$  is a submartingale with respect to  $\mathcal{H}_t$  under probability measure  $\mathbb{P}_{x_0}^{\tau, \theta}$ . It is crucial that  $Z_t^\gamma$  is a submartingale. [Kim \(2017\)](#) decomposes it into an  $\mathcal{H}_t$  martingale under  $\mathbb{P}_{x_0}^{\tau, \theta}$ ,

$$M_t^\gamma := \sum_{s=0}^t (\log \Lambda_s^\gamma - \mathbb{E}_{x_0}^{\tau, \theta}[\log \Lambda_s^\gamma | \mathcal{H}_{s-1}]),$$

and a predictable process,

$$A_t^\gamma := \sum_{s=0}^t \mathbb{E}_{x_0}^{\tau, \theta}[\log \Lambda_s^\gamma | \mathcal{H}_{s-1}].$$

We explain how the proof extends to our setting through [Assumption 1](#). The argument is twofold. The first requirement is that the increments of  $M_t^\gamma$  are bounded above and below by some  $d$ , i.e.,

$$|\log \Lambda_s^\gamma - \mathbb{E}_{x_0}^{\tau, \theta}[\log \Lambda_s^\gamma | \mathcal{H}_{s-1}]| \leq d, \tag{54}$$

for some  $d > 0$ . To see that (54) holds in our setting, note that by (37) and (38), i.e.,  $f^\gamma(r | x, u)$  and  $q^\gamma(y | x, u)$  are bounded away from 0, we have that  $|\log f^\gamma(r | x, u)| < \infty$  and  $|\log q^\gamma(y | x, u)| < \infty$ . This satisfies (54), which is needed for Azuma's inequality, a crucial step of Kim (2017)'s proof, to hold. The second requirement of the adaptation of the proof is

$$\sum_{s=0}^t \mathbb{E}_{x_0}^{\tau, \theta} [\log \Lambda_s^\gamma | \mathcal{H}_{s-1}] \geq \epsilon t, \quad (55)$$

i.e., (55) is an increasing predictable process. By (39), it follows that

$$\mathbb{E}_{x_0}^{\tau, \theta} [\log \Lambda_s^\gamma | \mathcal{H}_{s-1}] > \epsilon(x, u, \theta, \gamma) := \epsilon, \quad \forall s \leq t, \quad (56)$$

i.e., each increment of  $A_t^\gamma$  is strictly positive, which satisfies (55). The reader is referred to Kim (2017) for the details of why (56) holds. ■

*Proof of Lemma 7.1.* By the definition of almost-sure convergence, we have

$$\mathbb{P}_{x_0}^{\tau, \theta} \left( \lim_{t \rightarrow \infty} \pi_t(\theta | H_t) < 1 \right) = 1.$$

Therefore,

$$\begin{aligned} 1 &= \mathbb{P}_{x_0}^{\tau, \theta} \left( \lim_{t \rightarrow \infty} \pi_t(\theta | H_t) < 1 \right) = \mathbb{P}_{x_0}^{\tau, \theta} \left( \bigcup_{n=1}^{\infty} \left\{ \lim_{t \rightarrow \infty} \pi_t(\theta | H_t) < 1 - \frac{1}{n} \right\} \right) \\ &\leq \sum_{n=1}^{\infty} \mathbb{P}_{x_0}^{\tau, \theta} \left( \lim_{t \rightarrow \infty} \pi_t(\theta | H_t) < 1 - \frac{1}{n} \right). \end{aligned}$$

where the inequality is by Boole's inequality. Hence, Lemma 7.1 is verified. ■

## B Analysis of Assumption 1

Recall Assumption 1. Since Example 5.2 has a finite state space, (39) boils down to: For any  $x \in \mathcal{X}$ ,  $u \in \mathcal{U}$ , and any two distinct parameter value,  $\gamma \neq \theta \in \mathcal{P}$ , there exists a positive constant  $\epsilon(x, u, \theta, \gamma) > 0$  such that

$$\mathbb{K}(\nu_\theta^{x, u} | \nu_\gamma^{x, u}) \geq \epsilon(x, u, \theta, \gamma).$$

The above condition holds if and only if (by definition)

$$E_{f^\theta q^\theta} \left[ \log \left( \frac{d\nu_\theta^{x, u}}{d\nu_\gamma^{x, u}} \right) \right] = E_{f^\theta q^\theta} \left[ \log \left( \frac{f^\theta(\cdot | x, u) q^\theta(\cdot | x, u)}{f^\gamma(\cdot | x, u) q^\gamma(\cdot | x, u)} \right) \right] \geq \epsilon(x, u, \theta, \gamma),$$

where  $E_{f^\theta q^\theta}[\cdot | x, u]$  is defined in Section 3.2. For illustration purposes, we make the same assumption we had in Example 5.2; the  $\theta$ -MDP has only one state. This implies the transition kernel,

$q^\theta(\cdot \mid x, u)$ , is deterministic and we simplify the relative entropy expression. Due to the simpler version of the relative entropy, we utilize the expectation operator  $E_{f^\theta}[\cdot \mid x, u]$ , also defined in Section 3.2. It suffices to find a constant  $\epsilon(x, u, \theta, \gamma) > 0$ , for any  $x_A$  (in this case  $\mathcal{X} = \{x_0\}$  where  $x_0 = x_A$ ), any  $u \in \mathcal{U}$  (control 1 or 2),  $\theta = B$  and  $\gamma = A$ , such that

$$E_{f^\theta} \left[ \log \left( \frac{f^\theta(\cdot \mid x, u)}{f^\gamma(\cdot \mid x, u)} \right) \right] \geq \epsilon(x, u, \theta, \gamma).$$

We have,

$$\begin{aligned} E_{f^\theta} \left[ \log \left( \frac{\frac{1}{\sqrt{2\pi}0.1} e^{-\frac{(r-\mu^\theta)^2}{2(0.1)}} \mid x, u}{\frac{1}{\sqrt{2\pi}0.1} e^{-\frac{(r-\mu^\gamma)^2}{2(0.1)}} \mid x, u} \right) \right] &= E_{f^\theta} \left[ \log \left( \frac{e^{-(r-\mu^\theta)^2} \mid x, u}{e^{-(r-\mu^\gamma)^2} \mid x, u} \right) \right] \\ &= E_{f^\theta} \left[ \log \left( e^{-(r-\mu^\theta)^2 + (r-\mu^\gamma)^2} \mid x, u \right) \right], \end{aligned}$$

which can be simplified as

$$E_{f^\theta} [-(r - \mu^\theta)^2 + (r - \mu^\gamma)^2 \mid x, u].$$

**Case 1:** When the state is  $x_A$  and control 2 is picked by the TS policy,  $r \sim N(0.8, 0.1)$ , and

$$\begin{aligned} &E_{f^\theta} [-(r - 0.8)^2 + (r - 0.4)^2 \mid x_A, 2] \\ &= -Var[r \mid x_A, 2] + E_{f^\theta} [(r - 0.4)^2 \mid x_A, 2] \\ &= -0.1 + E_{f^\theta} [r^2 - 0.8r + 0.16 \mid x_A, 2] \\ &= -0.1 + E_{f^\theta} [r^2 \mid x_A, 2] - 0.8E_{f^\theta} [r \mid x_A, 2] + 0.16 \\ &= -0.1 + Var[r \mid x_A, 2] + (E_{f^\theta} [r \mid x_A, 2])^2 - 0.8E_{f^\theta} [r \mid x_A, 2] + 0.16 \\ &= -0.1 + 0.1 + 0.64 - 0.64 + 0.16 = 0.16. \end{aligned}$$

**Case 2:** When the state is  $x_A$  and control 1 is picked by the TS policy,  $r \sim N(0.3, 0.1)$ , and

$$\begin{aligned} &E_{f^\theta} [-(r - 0.3)^2 + (r - 0.5)^2 \mid x_A, 1] \\ &= -Var[r \mid x_A, 1] + E_{f^\theta} [(r - 0.5)^2 \mid x_A, 1] \\ &= -0.1 + E_{f^\theta} [r^2 - r + 0.25 \mid x_A, 1] \\ &= -0.1 + E_{f^\theta} [r^2 \mid x_A, 1] - E_{f^\theta} [r \mid x_A, 1] + 0.25 \\ &= -0.1 + Var[r \mid x_A, 1] + (E_{f^\theta} [r \mid x_A, 1])^2 - E_{f^\theta} [r \mid x_A, 1] + 0.25 \\ &= -0.1 + 0.1 + 0.09 - 0.3 + 0.25 = 0.04. \end{aligned}$$

As long as  $0 < \epsilon(x, u, \theta, \gamma) \leq 0.04$ , Assumption 1 holds for Example 5.2.

---

## References

- Agrawal S, Goyal N (2012) Analysis of thompson sampling for the multi-armed bandit problem. *Conference on Learning Theory*, 39–1.
- Banjević D, Kim MJ (2019) Thompson sampling for stochastic control: The continuous parameter case. *IEEE Transactions on Automatic Control* 64(10):4137–4152.
- Chung KL (2001) *A course in probability theory* (Academic press).
- Freedman DA (1963) On the asymptotic behavior of bayes' estimates in the discrete case. *The Annals of Mathematical Statistics* 1386–1403.
- Gopalan A, Mannor S (2015) Thompson sampling for learning parameterized markov decision processes. *Conference on Learning Theory*, 861–898.
- Hernández-Lerma O (2012) *Adaptive Markov control processes*, volume 79 (Springer Science & Business Media).
- Hernández-Lerma O, Lasserre JB (2012) *Discrete-time Markov control processes: basic optimality criteria*, volume 30 (Springer Science & Business Media).
- Kalkanli C, Ozgur A (2020) Asymptotic convergence of thompson sampling. *arXiv e-prints* arXiv–2011.
- Kearns M, Singh S (2002) Near-optimal reinforcement learning in polynomial time. *Machine learning* 49:209–232.
- Kim MJ (2017) Thompson sampling for stochastic control: The finite parameter case. *IEEE Transactions on Automatic Control* 62(12):6415–6422.
- Lai TL, Robbins H (1985) Asymptotically efficient adaptive allocation rules. *Advances in applied mathematics* 6(1):4–22.
- Leike J, Lattimore T, Orseau L, Hutter M (2016) Thompson sampling is asymptotically optimal in general environments. *arXiv preprint arXiv:1602.07905* .
- Puterman ML (2014) *Markov Decision Processes.: Discrete Stochastic Dynamic Programming* (John Wiley & Sons).
- Schäl M (1987) Estimation and control in discounted stochastic dynamic programming. *Stochastics: An International Journal of Probability and Stochastic Processes* 20(1):51–71.
- Shreve SE (1978) *Stochastic optimal control: The discrete time case* (Academic Press).
- Sutton RS, Barto AG (2018) *Reinforcement learning: An introduction* (MIT press).
- Thompson WR (1933) On the likelihood that one unknown probability exceeds another in view of the evidence of two samples. *Biometrika* 25(3/4):285–294.